

## ⇒ Research Article



# Effect of Pre-conditioning of Endurance Training with Different Intensities on Soleus Muscle Atrophy in a Period of Inactivity: The Role of PGC-1 $\alpha$ 4 Gene

Aram Asadi Karam<sup>1</sup>, Javad Nemati<sup>1</sup>, Majid Asadi-Shekaari<sup>2\*</sup>, Farhad Daryanoosh<sup>1</sup>, Mohammad Hemmatinfar<sup>1</sup>

<sup>1</sup>Department of Exercise Physiology, Faculty of Psychology and Educational Science, Shiraz University, Shiraz, Iran

<sup>2</sup>Neuroscience Research Center, Neuropharmacology Institute, Kerman University of Medical Sciences, Kerman, Iran

## Abstract

**Background:** This study aimed to measure PGC-1 $\alpha$ 4 soleus muscle and to investigate the effect of endurance training pre-conditioning with different intensities on the atrophy response of the soleus muscle to a period of inactivity.

**Methods:** In this experimental study, 24 male Wistar rats were randomly divided into equal groups of control (C), control inactivity (CI), high-intensity endurance training group (HE) (treadmill speed 30 km/h), and low-intensity endurance training (LE) (treadmill's speed: 10 km/h). After two weeks of familiarization, the endurance training group ran on a treadmill for two weeks (five sessions each week). The animal's lower limbs were then immobilized for seven days. Then the soleus muscle was extracted, and after weighing, the expression of the PGC-1 $\alpha$ 4 gene was measured using the real-time polymerase chain reaction (real-time PCR) technique. Data were analyzed using SPSS software, version 24.

**Results:** The expression of the PGC-1 $\alpha$ 4 gene was significantly higher in the HE group than the CI group. However, compared to the C group, all groups with inactivity intervention showed significantly lower PGC-1 $\alpha$ 4 gene levels. The ratio of muscle mass to body weight in the C group was significantly higher than the LE and CI groups, and higher in the HE group than the CI group.

**Conclusion:** Endurance training seems to be able to reduce the destructive effects of inactive atrophy. The higher intensity of these exercises was more effective, which was associated with increased expression of the PGC-1 $\alpha$ 4 gene.

**Keywords:** Atrophy, Endurance exercise, Inactivity, PGC-1 $\alpha$ 4

## \*Correspondence to

Majid Asadi-Shekaari,  
Email: Majidasadi@kmu.ac.ir



Received November 9, 2020, Accepted: September 25, 2021, Published Online: September 29, 2021

## Background

Muscle atrophy significantly reduces the ability to perform physical tasks leading to decreased independence and increased mortality (1). Skeletal muscle atrophy occurs in pathological (such as cancer and diabetes) and non-pathological (such as inactivity and muscle disuse) conditions. It has been suggested that the mentioned conditions lead to loss of muscle protein, muscle fibrous atrophy, and functional impairment (2). Atrophy is caused by muscle disuse identified by a decrease in the size, weight, and function of skeletal muscle (3). In muscle disuse, because of the imbalance in synthesis and degradation of the protein, a rapid decrease in muscle size has been observed (4). Anabolic pathways include the mechanical stretching pathways (5), the mammalian target of rapamycin (mTOR) (6), the Wnt/ $\beta$ -catenin pathway, and ribosomal biogenesis (7), the beta-adrenergic receptor pathway (8), plus some unknown and emerging pathways such as nitric oxide (NO) (9), peroxisome proliferator-activated receptor-gamma coactivator-1 alpha (PGC-1 $\alpha$ ) (10), and microRNAs (4).

PGC-1 $\alpha$ 4, one of the isoforms of PGC-1 $\alpha$ , promotes muscle growth by activating IGF-1 and inhibiting myostatin (11). Although PGC-1 $\alpha$ 4 expression decreases in atrophic conditions, its changes in inactivity have not been well established (12). However, the expression of PGC-1 $\alpha$ 4 in the skeletal muscle of rat causes resistance to cancer and muscle atrophy induced by disuse (13).

Research on the preventive factors in muscle atrophy is scarce (14). Some studies have investigated the effective nutritional and pharmacological approaches in the prevention of atrophy. However, physical activity as a non-pharmacological strategy is effective (15). Physical activity before the period of inactivity or sedentary state can reduce the rate of muscle atrophy (16); however, the factors and mechanisms involved are not clear.

Several studies have shown the role of aerobic exercise pre-conditioning in reducing atrophy after inactivity. Decreasing the expression of PGC-1 $\alpha$  gene and protein induced by inactivity, in mice with five weeks of endurance training was lower than not trained mice (17). Also, PGC-1 $\alpha$ 4 expression was significantly increased in

response to an eight-week exercise program that included resistance training (10). This result was obtained following endurance training as well (18).

Despite the proven positive effects of intensity on muscle adaptation through factors such as metabolic stress, accumulation of metabolites such as lactate, mineral phosphate (Pi), and hydrogen ions (H) that have a significant effect on hormone release, hypoxia, cell swelling, and production of reactive oxygen species which can initiate anabolic pathways for muscle growth and adaptation to energy metabolism (19), the effect of different intensities of endurance training on the role of new factors such as PGC-1 $\alpha$ 4 in anabolic and catabolic pathways involved in the pretraining of skeletal muscle for confronting atrophy induced by inactivity or not using the limb is not well defined (19).

We aimed to measure PGC-1 $\alpha$ 4 of muscle tissue and evaluate the effect of pre-conditioning of endurance training with different intensities on the response of the soleus muscle atrophy in a period of inactivity in male rats.

## Material and Methods

### Techniques and Samples

Twenty-four Wistar rats aged eight weeks (weight: 180  $\pm$  20 g) were used. After 2 weeks of acclimatization in the animal room at a temperature of 22  $\pm$  3°C, 40-60% humidity, 24 sleep-wake cycle (12 hours of light and 12 hours of darkness), and free access to water and food, they were randomly divided into four groups of six: the control (C); control of inactivity (CI); low-intensity endurance training (LE); high-intensity endurance training (HE). Induction of immobility was performed in LE and HE groups after two weeks of training (five days a week) and in the CI group performed without training.

### Exercise Group Protocol

When groups underwent a familiarization period and the training time progressively reached 30 minutes in all groups, the LE and HE groups trained at 10 and 30 m/min, respectively. To match the stress on the treadmill, group C experienced walking on a treadmill at 5 m/s for 15 minutes (20, 21).

### Induction of Immobility by Hanging Lower Limb

After anesthetizing the rat, a wire was passed through the tail, and a ring made from wire then connected it to the screw that had hung from the lid of the container with a clamp. This period lasted for seven days (22).

### Isolation and Tissue Preparation

After deep anesthetizing with ketamine (75 mg/kg) and xylazine (10 mg/kg), blood samples were taken from the heart, and the soleus muscles were removed. The muscles were rinsed in saline and weighed to determine

hypertrophy/atrophy and were frozen (-80°C) until further tests.

## Evaluation of PGC-1 $\alpha$ 4 Gene Expression in Soleus Muscle Tissue

### RNA Extraction

This step was done by the RNA extraction kit (CinnaGen Inc., Tehran, Iran). For achieving this goal, tissues (50 mg) were lysed by trizol solution and completely homogenized with a tissue homogenizer and then the aqueous phase was separated using 0.25 mL of chloroform. Extracted RNA was washed and dehydrated by 1ml cold ethanol (70%). Then sterile water was added (5.1  $\mu$ L/mg tissue). The quantity and quality of the obtained RNA were checked by measuring the ratio of the optical density of 260/280 nm using Nanodrop™ spectrophotometer (Nanodrop; Thermo Fisher Scientific, Wilmington, DE, USA).

### cDNA Synthesis

cDNA Synthesis kit (Thermo Fisher Scientific, Inc., Waltham, MA, USA) was used for cDNA synthesis. For each sample, three steps of cDNA synthesis were performed. Thus, at first, 8  $\mu$ L of the extracted RNA was mixed with 0.8  $\mu$ L of DNase I enzyme, 2  $\mu$ L of 10x buffers, and DEPC water then brought the mixture up to a total volume of 20  $\mu$ L. The final product was gently mixed without vortexing and then incubated in the thermocycler with the following program: 5 min at 55°C, 15 minutes at 25 °C, 30 minutes at 42 °C (cDNA synthesis step by RT enzyme), 5 minutes at 95°C (to inactivation of RT enzyme).

After completion of the thermocycler steps, 280  $\mu$ L of injection water was added and stored at -20°C for use in quantitative polymerase chain reaction (qPCR). Also, for each cDNA sample, a positive control sample with a b2m primer was prepared as an internal control and test for the presence of cDNA. The samples were mixed slowly and without vortex and placed in a real-time PCR set (StepOne, Applied Biosystems, USA) with the following program: 10 minutes at 95°C (initial denaturation), 10 seconds at 95°C (denaturation), 15 seconds at 60°C (primers binding), and 20 seconds at 72°C (extension). The reaction was repeated from the second stage onwards for 40 cycles. Threshold cycles (Cts) related to the reactions were extracted by real-time PCR software and finally, the mean three times Cts was recorded.

### Quantification of Target Gene Expression Levels

To calculate gene expression levels, the 2<sup>- $\Delta\Delta$ CT</sup> ("delta-delta Ct") method was used.

### Statically Analysis

All results were expressed in different groups as mean  $\pm$  SD. The data were analyzed using SPSS software, version 21. After checking the normality of the data

with Kolmogorov-Smirnov test, One-way analysis of variance (ANOVA) and Tukey's post hoc tests were used for statistical analysis and comparison between groups. Pearson's correlation test was used to evaluate the relationship between PGC-1 $\alpha$ 4 gene expression levels and soleus muscle hypertrophy. In all cases,  $P < 0.05$  was considered statistically significant.

## Results

### Mean Body Weight

The mean body weight at the beginning and end of the study in all groups is shown in Table 1. There was a significant difference in the final weight between the four groups ( $F_{3,23} = 13.02, P < 0.01$ ). The final weight in group C was significantly higher than groups CI ( $P = 0.000$ ), LE ( $P = 0.001$ ), and HE ( $P = 0.041$ ). Also, the final weight of the HE group was significantly higher than the CI group ( $P = 0.033$ ).

### Mean Soleus Muscle Weight and its Ratio to Body Weight

The mean weight of the soleus muscle and its ratio to body weight, as an important indicator of hypertrophy and muscle atrophy in different groups, is shown in Table 2. There was a statistically significant difference between the four groups in this regard ( $F_{3,23} = 10.1, P < 0.01$ ). Tukey's post-hoc test showed that this index was significantly higher in the HE group compared with the CI group. On the other hand, this index was significantly lower than the C group in LE ( $P = 0.006$ ) and CI ( $P < 0.001$ ) groups. This ratio was lower in the HE group compared with the C group ( $P = 0.215$ ).

### PGC-1 $\alpha$ 4 Levels of Soleus Muscle Tissue

There was a significant difference in the PGC-1 $\alpha$ 4 level

**Table 1.** The Mean  $\pm$  SD Weight of the Groups (n=6) at the Beginning and End of the Exercise

Groups	Initial Weight (g)	Final Weight (g)
C	183.20 $\pm$ 3.20	311.33 $\pm$ 7.71*
HE	182.44 $\pm$ 2.67	297.50 $\pm$ 9.73 <sup>#</sup>
LE	185.17 $\pm$ 4.42	289.33 $\pm$ 7.71
CI	184.27 $\pm$ 2.24	283.16 $\pm$ 7.71

\*  $P < 0.05$  as compared to other groups; #  $P < 0.05$  as compared to CI group. Control (C); Control of Inactivity (CI); Low-intensity endurance training (LE); High-intensity endurance training (HE).

**Table 2.** The Mean  $\pm$  SD Weight of the Soleus Muscle in Different Groups (n=6) and its Ratio to Bodyweight

Groups	Weight of the Soleus Muscle (mg)	Sol /BW (mg/g)
C	150.83 $\pm$ 4.66	0.48 $\pm$ 0.02*
HE	137.33 $\pm$ 4.71	0.46 $\pm$ 0.01 <sup>#</sup>
LE	127.83 $\pm$ 5.52	0.44 $\pm$ 0.02
CI	120.50 $\pm$ 1.87	0.42 $\pm$ 0.02

\*  $P < 0.05$  as compared to CI and LE groups; #  $P < 0.05$  as compared to CI group. Control (C); Control of inactivity (CI); Low-intensity endurance training (LE); High-intensity endurance training (HE); soleus muscle's weight to body weight (Sol/BW)

in soleus muscle tissue in different groups ( $F_{3,23} = 27.43, P < 0.01$ ). The PGC-1 $\alpha$ 4 level in the groups that had inactivity intervention (HE, LE, and CI) was significantly lower than the C group ( $P < 0.01$ ). In the HE group, the PGC-1 $\alpha$ 4 level was significantly higher than the CI group (without exercise,  $P < 0.01$ ). Also, the PGC-1 $\alpha$ 4 level in the LE group was higher than the CI group, but this difference was not statistically significant (Figure 1).

### Relationship Between PGC-1 $\alpha$ 4 Gene Expression and Soleus Muscle Hypertrophy

The results of Pearson's correlation test did not show a significant relationship between PGC-1 $\alpha$ 4 gene expression values and soleus muscle hypertrophy in the three groups (Table 3).

## Discussion

This study showed the beneficial effects of pre-conditioning with a period of endurance exercise training in reducing the harmful effects of seven days of non-training on the soleus muscle in rats. This period of pre-conditioning greatly reduced the loss of muscle mass in the endurance exercise groups (HE and LE) compared with the control group. Moreover, the expression levels of proteogenesis pathways in endurance exercise groups were higher than in the control group. Accordingly, these findings reinforce the hypothesis that endurance exercise pre-conditioning reduces harmful changes in the soleus muscle.


We found that endurance training reduced atrophy after inactivity. This result was similar to another study showing less reduction in muscle hypertrophy and expression of the PGC-1 $\alpha$  gene and protein in the group that had five weeks of endurance exercise before inactivity (17). Other researchers reported an increase in PGC-1 $\alpha$ 4 gene expression in response to the exercise and resistance training program (18, 23).

Concerning PGC-1 $\alpha$ 4 levels and hypertrophy index, after the inactivity period, the endurance training groups (HE and LE groups) and the CI group had lower results compared to the control group (without intervention), but the soleus muscle hypertrophy index was significantly increased in the HE group compared with the CI group by 9.5%. This increase was also seen in the LE group (4.7%), but was not statistically significant. This increase can probably be related to the higher intensity of exercise activity in the HE group. On the other hand, similar to the results of the hypertrophy index, the level of PGC-1 $\alpha$ 4 in

**Table 3.** Correlation Between PGC-1 $\alpha$ 4 Gene Expression Level and Soleus Muscle Hypertrophy in Different Groups (n=6)

Correlation of Variables	HE	LE	CI
Gene expression level and muscle hypertrophy	$r = 0.23$ ( $P = 0.65$ )	$r = -0.68$ ( $P = 0.13$ )	$r = -0.44$ ( $P = 0.37$ )

Control of Inactivity (CI); Low-intensity endurance training (LE); High-intensity endurance training (HE).



**Figure 1.** Mean of the PGC1- $\alpha$ 4 level of soleus muscle tissue in different groups  
\*  $P < 0.05$  as compared to C group; #  $P < 0.05$  as compared to HE group  
Control (C); Control of Inactivity (CI); Low-intensity endurance training (LE); High-intensity endurance training (HE)

the groups with inactivity intervention was significantly lower than the control group (without exercise and inactivity). However, the PGC-1 $\alpha$ 4 gene expression level in the soleus muscle of the HE group was significantly higher than the CI group (without exercise intervention). Also, in the period of inactivity, the effects of inactivity on soleus muscle mass have been shown. The intensity of exercise increases the expression of the PGC-1 $\alpha$ 4 gene, which can play an important role in the process of hypertrophy and the prevention of atrophy. The proteogenesis pathway of PGC-1 $\alpha$ 4 is likely to be further stimulated by increasing the intensity of endurance training, affecting slow-twitch fibers such as the soleus muscle. Although most research have shown an association between PGC-1 $\alpha$ 4 and muscle hypertrophy with resistance training, studies have shown that PGC-1 $\alpha$ 4 also increases with endurance exercise (18, 24) and hypoxia (25). In general, PGC-1 $\alpha$ 4 promotes muscle growth by activating IGF-1 and inhibiting myostatin and prevents the processes of muscle atrophy because of inactivity or underload and disease. Endurance exercise may suppress muscle atrophy mechanisms by reducing inflammatory cytokines in the long-term in muscles, reducing free oxygen species, and increasing PGC-1 $\alpha$  gene expression. In the period of lack of training and weightlessness, the expression of proteins of these genes and the mentioned changes remain at a high level. On the other hand, direct stimulation of IGF-1 activates the growth pathways related to the PI3K-AKT-mTOR cell cascade and prevents atrophy in the long-term, and promotes protein production. However, according to the results of the present study, inactivity, even when the rats trained, was able to significantly reduce the levels of PGC-1 $\alpha$ 4 and soleus hypertrophy index compared with the control group. Muscle growth and expression of PGC-1 $\alpha$ 4 gene in the soleus muscle during exercise were much higher than the control group, for this reason, with inactivity; a larger reduction slope was created. It seems that the higher the baseline level of a physiological factor, the more significant the effect of reduction interventions.

It seems that increasing the intensity of endurance training can further stimulate factors such as higher metabolic stress, accumulation of stimulant metabolites such as lactate, mineral phosphate (Pi), and hydrogen ions (H) that affect hormonal release, hypoxia, cell swelling, and the production of reactive oxygen species and further stimulate anabolic pathways for muscle growth and adaptation to energy metabolism (19). One of the most important growth hormone stimuli involved in skeletal muscle hypertrophy is lactate accumulation, which can stimulate NO to release more growth hormone in addition to NO stimulation.

Higher expression of PGC-1 $\alpha$ 4 gene in the period of inactivity in HE and LE groups compared to the CI group can also be attributed to adaptations related to antioxidant defenses because in the period of inactivity, by reducing the adaptation of muscle fibers, free radicals increase and destroy muscle fibers, especially slow-twitch fibers, leading to muscle atrophy.

The positive correlation between PGC-1 $\alpha$ 4 gene expression level and soleus muscle mass in the HE group indicates that high-intensity training has been able to prevent muscle mass loss in the inactivity period by higher pre-conditioning changes than the LE group. Because of the lack of measurement of muscle mass at the end of the last training step and before the start of the inactivity period, it is not clear whether endurance training with higher intensity caused more hypertrophy in the soleus muscle or it had prevented the further reduction of muscle mass during the inactivity period. The reason for the lack of a positive relationship between muscle hypertrophy index and expression level of the PGC-1 $\alpha$ 4 gene in the soleus muscle of the HE group may be related to the sample size of the study (six heads). By measuring the expression level of the two main targets of the PGC-1 $\alpha$ 4 gene, IGF-1, and myostatin, it may be possible to talk more accurately about the effects of endurance exercise as an important factor in pre-preparation for inactivity or injury.

## Conclusion

The results of this study showed that endurance exercise as a non-pharmacological intervention can create beneficial effects and adaptations in the soleus muscle, which can greatly prevent destruction, atrophy, and reduced muscle force in the period of inactivity through pre-conditioning. The higher the intensity of this type of exercise, the greater the pre-conditioning level. One of the most important indicators is the PGC-1 $\alpha$ 4 gene expression level, but to determine the more accurate and clear paths, more studies are needed in the upstream and downstream paths of this indicator.

## Authors' Contribution

All those who have mentioned their names as authors in this article confirm that they have contributed to the preparation of content,

concept, design, analysis, and interpretation of data as well as drafting the manuscript. All authors are responsible for writing this article.

#### Ethical Approval

Procedures adopted in our study were under the ethical guidelines of the Ethics Committee of Kerman University of Medical Sciences for laboratory animals (code: IR.KMU.EC.1399,368) that were compatible with the Helsinki Declaration on the Protection of Vertebrates for Laboratory and Other Scientific Purposes.

#### Conflict of Interests

None declared.

#### Funding/Support

The authors have received financial support from Kerman Neuroscience Research Center for this study.

#### Acknowledgments

The authors would like to thank the staff of Kerman Neuroscience Research Center for their kind support.

#### References

1. Powers SK, Lynch GS, Murphy KT, Reid MB, Zijdewind I. Disease-induced skeletal muscle atrophy and fatigue. *Med Sci Sports Exerc.* 2016;48(11):2307-19. doi: 10.1249/mss.0000000000000975.
2. Theilen NT, Jeremic N, Weber GJ, Tyagi SC. Exercise preconditioning diminishes skeletal muscle atrophy after hindlimb suspension in mice. *J Appl Physiol* (1985). 2018;125(4):999-1010. doi: 10.1152/japplphysiol.00137.2018.
3. Theilen NT, Kunkel GH, Tyagi SC. The role of exercise and TFAM in preventing skeletal muscle atrophy. *J Cell Physiol.* 2017;232(9):2348-58. doi: 10.1002/jcp.25737.
4. McCarthy JJ, Murach KA. Anabolic and catabolic signaling pathways that regulate skeletal muscle mass. In: Bagchi D, Nair S, Sen CK, eds. *Nutrition and Enhanced Sports Performance.* 2nd ed. Academic Press; 2019. p. 275-90. doi: 10.1016/b978-0-12-813922-6.00024-2.
5. Crossland H, Kazi AA, Lang CH, Timmons JA, Pierre P, Wilkinson DJ, et al. Focal adhesion kinase is required for IGF-I-mediated growth of skeletal muscle cells via a TSC2/mTOR/S6K1-associated pathway. *Am J Physiol Endocrinol Metab.* 2013;305(2):E183-93. doi: 10.1152/ajpendo.00541.2012.
6. Laplante M, Sabatini DM. mTOR signaling in growth control and disease. *Cell.* 2012;149(2):274-93. doi: 10.1016/j.cell.2012.03.017.
7. Rommel C, Bodine SC, Clarke BA, Rossman R, Nunez L, Stitt TN, et al. Mediation of IGF-1-induced skeletal myotube hypertrophy by PI(3)K/Akt/mTOR and PI(3)K/Akt/GSK3 pathways. *Nat Cell Biol.* 2001;3(11):1009-13. doi: 10.1038/ncb1101-1009.
8. Yamamoto DL, Hutchinson DS, Bengtsson T. Beta(2)-adrenergic activation increases glycogen synthesis in L6 skeletal muscle cells through a signalling pathway independent of cyclic AMP. *Diabetologia.* 2007;50(1):158-67. doi: 10.1007/s00125-006-0484-0.
9. Sellman JE, DeRuisseau KC, Betters JL, Lira VA, Soltow QA, Selsby JT, et al. In vivo inhibition of nitric oxide synthase impairs upregulation of contractile protein mRNA in overloaded plantaris muscle. *J Appl Physiol* (1985). 2006;100(1):258-65. doi: 10.1152/japplphysiol.00936.2005.
10. Ruas JL, White JP, Rao RR, Kleiner S, Brannan KT, Harrison BC, et al. A PGC-1 $\alpha$  isoform induced by resistance training regulates skeletal muscle hypertrophy. *Cell.* 2012;151(6):1319-31. doi: 10.1016/j.cell.2012.10.050.
11. Schiaffino S, Dyar KA, Cicilioti S, Blaauw B, Sandri M. Mechanisms regulating skeletal muscle growth and atrophy. *FEBS J.* 2013;280(17):4294-314. doi: 10.1111/febs.12253.
12. Bowen TS, Schuler G, Adams V. Skeletal muscle wasting in cachexia and sarcopenia: molecular pathophysiology and impact of exercise training. *J Cachexia Sarcopenia Muscle.* 2015;6(3):197-207. doi: 10.1002/jcsm.12043.
13. Martínez-Redondo V, Pettersson AT, Ruas JL. The hitchhiker's guide to PGC-1 $\alpha$  isoform structure and biological functions. *Diabetologia.* 2015;58(9):1969-77. doi: 10.1007/s00125-015-3671-z.
14. Smuder AJ, Kavazis AN, Min K, Powers SK. Exercise protects against doxorubicin-induced markers of autophagy signaling in skeletal muscle. *J Appl Physiol* (1985). 2011;111(4):1190-8. doi: 10.1152/japplphysiol.00429.2011.
15. Powers SK. Can antioxidants protect against disuse muscle atrophy? *Sports Med.* 2014;44 Suppl 2:S155-65. doi: 10.1007/s40279-014-0255-x.
16. Adams GR, Haddad F, Bodell PW, Tran PD, Baldwin KM. Combined isometric, concentric, and eccentric resistance exercise prevents unloading-induced muscle atrophy in rats. *J Appl Physiol* (1985). 2007;103(5):1644-54. doi: 10.1152/japplphysiol.00669.2007.
17. Aoi W, Naito Y, Mizushima K, Takanami Y, Kawai Y, Ichikawa H, et al. The microRNA miR-696 regulates PGC-1 $\alpha$  in mouse skeletal muscle in response to physical activity. *Am J Physiol Endocrinol Metab.* 2010;298(4):E799-806. doi: 10.1152/ajpendo.00448.2009.
18. Lundberg TR, Fernandez-Gonzalo R, Norrbom J, Fischer H, Tesch PA, Gustafsson T. Truncated splice variant PGC-1 $\alpha$ 4 is not associated with exercise-induced human muscle hypertrophy. *Acta Physiol (Oxf).* 2014;212(2):142-51. doi: 10.1111/apha.12310.
19. de Freitas MC, Gerosa-Neto J, Zanchi NE, Lira FS, Rossi FE. Role of metabolic stress for enhancing muscle adaptations: Practical applications. *World J Methodol.* 2017;7(2):46-54. doi: 10.5662/wjm.v7.i2.46.
20. Kang C, Ji LL. Muscle immobilization and remobilization downregulates PGC-1 $\alpha$  signaling and the mitochondrial biogenesis pathway. *J Appl Physiol* (1985). 2013;115(11):1618-25. doi: 10.1152/japplphysiol.01354.2012.
21. Kang C, Ji LL. Muscle immobilization and remobilization downregulates PGC-1 $\alpha$  signaling and the mitochondrial biogenesis pathway. *J Appl Physiol* (1985). 2013;115(11):1618-25. doi: 10.1152/japplphysiol.01354.2012.
22. Ferreira JA, Crissey JM, Brown M. An alternant method to the traditional NASA hindlimb unloading model in mice. *J Vis Exp.* 2011(49):2467. doi: 10.3791/2467.
23. Rossignol DA, Frye RE. A review of research trends in physiological abnormalities in autism spectrum disorders: immune dysregulation, inflammation, oxidative stress, mitochondrial dysfunction and environmental toxicant exposures. *Mol Psychiatry.* 2012;17(4):389-401. doi: 10.1038/mp.2011.165.
24. Ydfors M, Fischer H, Mascher H, Blomstrand E, Norrbom J, Gustafsson T. The truncated splice variants, NT-PGC-1 $\alpha$  and PGC-1 $\alpha$ 4, increase with both endurance and resistance exercise in human skeletal muscle. *Physiol Rep.* 2013;1(6):e00140. doi: 10.1002/phy2.140.
25. Thom R, Rowe GC, Jang C, Safdar A, Arany Z. Hypoxic induction of vascular endothelial growth factor (VEGF) and angiogenesis in muscle by truncated peroxisome proliferator-activated receptor  $\gamma$  coactivator PGC-1 $\alpha$ . *J Biol Chem.* 2014;289(13):8810-7. doi: 10.1074/jbc.M114.554394.