

Serum concentration of myostatin, insulin and glucose response to continuous and discontinuous exercises in overweight menopausal and menstrual women

Masomeh Hannani¹ Abbas Ali Gaeini² Ramin Shabani³ Mohammad Hemmatinafar⁴

PhD Student of Exercise Physiology¹, Tehran University Kish International Camps, Kish, Iran. Professor Department of Exercise Physiology², Tehran University, Tehran, Iran. Associate Professor Department of Exercise Physiology³, Islamic Azad University, Rasht Branch, Rasht, Iran. Assistant Professor Department of Exercise Physiology⁴, Shiraz University, Shiraz, Iran.

(Received 29 Jan, 2015

Accepted 8 Apr, 2016)

Original Article

Abstract

Introduction: Myostatin regulates skeletal muscle mass and adipogenesis, but serum response to continuous and discontinuous exercises training it has not been studied in menopause and menstrual groups of women. The purpose of this study was to investigate the serum response of myostatin, insulin and glucose to use four different aerobic exercises training in menopause and menstrual groups of women.

Methods: Seventeen overweight women (BMI>30 and 40-50 years old) in Bandar abbas voluntarily participated in this study and randomly divided into two groups: menopausal (n=8) and menstrual (n=9) groups. Myostatin, insulin and glucose were measured before and after four different exercise sessions including continuous and discontinuous aerobic training in water and on treadmill. Data were analyzed by paired t-test and Analysis of covariance.

Results: Myostatin levels were significantly decreased after continuous and discontinuous training in water, and discontinuous training on treadmill in menopause group ($P<0.05$). Meanwhile, myostatin level showed significant decrease in all four different conditions of training in menstrual group ($P<0.05$). Insulin levels were significantly decreased after continuous and discontinuous training in water, and continuous training on treadmill in both menstrual and menopause groups ($P<0.05$). Glucose levels were significantly decreased after continuous and discontinuous training in water, and continuous training on treadmill in menopause group ($P<0.05$). In menstrual group, glucose levels were significantly decreased after continuous and discontinuous training on treadmill, and discontinuous training in water ($P<0.05$).

Conclusion: The lowest myostatin response was seen in discontinuous training in water in both menopause and menstrual groups, while the highest myostatin response observed in menopause group who committed continuous training on treadmill. This study suggested that discontinuous than continuous exercise, and exercise in the water than on treadmill, Lead to better response in variables are listed.

Correspondence:
Abbas Ali Gaeini, PhD.
Exercise Physiology
Department, Faculty of
Physical Education & Sport
Sciences., Tehran University.
Tehran, Iran
Tel: +98 9123351872
Email:
aagaieini@yahoo.com

Key words: Exercise, Overweight, Menopause, Myostation

Citation: Hannani M, Gaeini AA, Shabani R, Hemmatinafar M. Serum concentration of myostatin, insulin and glucose response to continuous and discontinuous exercises in overweight menopausal and menstrual women. Hormozgan Medical Journal 2016;20(2):80-87.

Introduction:

Aging is associated with a wide range of health deficiency. In this regard, menopause is a status in the aging process of women, which marks the cessation of menstrual cycle (1). Also, It is associated with dramatic reduction in circulating estrogen and progesterone concentrations in body which can result in reduced insulin sensitivity and consequently elevated risk of diabetes, and adipose tissue enhancement predominantly in the abdominal region and subsequently an increase in cardiovascular diseases, hyperglycemia and hyperinsulinemia (2-4). Available literatures support this notion that aerobic activities are useful for prevention or treatment of age-associated weight and fat gain, and therefore obesity-associated metabolic changes and diseases (5,6).

Also, sarcopenia, reduced muscle mass by aging that may be induced by elevated expression of myostatin, can lead to the enhanced risk of cardiac, pulmonary, and metabolic diseases (7). Myostatin belongs to the transforming growth factor- β super family and is a growth factor protein that limits muscle tissue growth (7). The myostatin protein is produced primarily in skeletal muscle cells, circulates in the blood and lymph and acts on muscle tissue, apparently by slowing down the development of muscle stem cells (8). Recent evidence imply on myostatin reduction in response to muscle exercise e.g. acute and short-term swimming training; chronic wheel running; and muscle loading through treadmill running or isometric resistance training after atrophy induced by hindlimb unloading (8).

So, according to all above mentioned reasons, successful physical activities to reduce overweight and the dire consequences associated with aging are of utmost importance in both prevention and treatment of obesity and metabolic syndrome (9).

The main objective of this study was to examine myostatin, insulin and glucose responses to continuous and discontinuous aerobic training on treadmill and in water in overweight menopausal and menstrual women. It is hypothesized that following aerobic training in water and on treadmill, serum myostatin, insulin and glucose levels would be decreased.

Methods:

This is an interventional study. Nineteen overweight individuals, who had no previous history of routine exercise training, enrolled in our investigation. Two individuals did not complete the study because of either time constraints or medical issues unrelated to study participation. The 17 individuals (8 menopausal women and 9 menstrual women) who completed the study were more than 40 years old and had body mass index more than 30 kg/m². Informed consent was obtained from all participants, together with the manner in which informed consent was obtained.

The training protocols were designed to provide 70-80% maximum heart rate, $H.R.Max = 208 - (0.7 \times age)$, for 30 minutes. Subjects trained 1 time per week across 4-weeks. Training sessions included two types of discontinuous and continuous programs. Participants in discontinuous group performed 3 sets of 10 minutes repetitions with 5 minutes resting time at every session, while participants in continuous group performed a continuous exercise in a 30-minutes session.

Training protocols involved four different programs with one week interval between each one. Training sessions started with a continuous aerobic training in water in the first week, and followed by a discontinuous aerobic training on treadmill in the second week, a discontinuous aerobic training in water in the third week and a continuous aerobic training on treadmill in the fourth week.

Training program in water began with 5-10 minutes stretching and warm-up phase then followed by aerobic exercise in water and ends up with 5 minutes cool-down phase. Whilst, training program on treadmill started with 5-10 minutes slow walking on treadmill for warming, and then followed to reach approximately 70-80% maximum heart rate and ends up with 5 minutes cool-down.

All stages the exercise intervention was performed in a sports complex in the city of Bandar Abbas.

Blood samples were drawn before and after each training sessions. Blood samples were drawn from a superficial arm vein and were centrifuged at 1000 rpm for 15 min within 30 minutes of collection for obtaining serum. Serum samples were then aliquated into separate tubes and stored at -20 ° C till analysis.

Body composition Analysis ioi 353 was used for measurement of height, weight and Body Mass Index (BMI), which calculated by dividing body weight by square of height in meter, of participants.

Serum myostatin concentrations were assessed using a competitive enzyme linked immunosorbent assay (ELISA) method (Human Myostatin ELISA Kit, Cusabio Biotech, Chin). Initially, 50 μ l of standard or sample per well were added. Then, 50 μ l of HRP-conjugate was added to each well. Plate was incubated for 1 hour at 37° C. The liquid of each well was aspirated and washed for three times. After the last wash, any remaining liquid was removed by aspirating. Then, substrates were added to each well and incubated for 15 minutes at 37°C. Finally, stop solution was added and optical density was read using a microplate reader at 450 nm (10).

Insulin serum levels were measured by ELISA method (Mercodia insulin ELISA; Mercodia AB, Uppsala, Sweden). At first, 25 μ l each of Calibrators and samples were poured into wells. Then, 100 μ l of enzyme conjugate was added to each well. Plate was incubated on a plate shaker for 1 hour at room temperature. After washing, 200 μ l Substrate TMB was added into each well and incubated for 15 minutes at room temperature (18–25°C). After adding Stop Solution, optical density was read at 450nm (11).

For determination of glucose levels, 1ml of reagent was added to all tubes. Then, 10 μ l of standard or samples were added to tubes (one tube was filled with 10 μ l of distilled water as blank) and incubated for 20 minutes at room temperature. Finally, absorbance against the blank was read at 546 nm (12,13).

For comparison of myostatin, insulin and glucose response before and after four different training programs in two menopause and menstrual women groups, t-tests analysis was applied. Analysis of covariance was used to examine

differences between groups after statistically removing the effect of selected covariates. Data were analyzed using SPSS software version 19.0. Data are presented as means SEM. Probability $P < 0.05$ is statistically significant.

Results:

Body composition is a reliable indicator to find possible health problems derived from the amount of body-composing constituents and imbalance between them. In this regard, the mean (\pm standard deviation) of body composition and age of menopause and menstrual women are indicated in Table 1.

Table 1. The mean and standard deviation of participants' characteristics are indicated

Groups	Age	Height (cm)	Weight (kg)	BMI
Menopause	51.37 \pm 6.04	158.77 \pm 4.72	82.72 \pm 11.79	33.16 \pm 3.30
Menstrual	47.37 \pm 2.91	159.33 \pm 5	88.95 \pm 10.93	34.98 \pm 3.46

In menopause group, myostatin levels in continuous / water, discontinuous / treadmill, discontinuous/water and continuous treadmill groups were decreased -1.36 ± 0.63 ($P=0.000$) from 9.17 ± 4.38 to 7.81 ± 4.21 , $-1.65 \pm .92$ ($P=0.000$) from 9.11 ± 4.43 to 7.46 ± 4.32 , -2.075 ± 1.64 ($P=0.000$) from 9.69 ± 4.60 to 7.61 ± 3.72 , and -1.21 ± 2.48 ($P= 0.210$) from 9.07 ± 4.14 to 7.86 ± 3.53 respectively.

In menstrual group, myostatin levels in continuous / water, discontinuous treadmill, discontinuous / water and continuous treadmill groups were decreased -1.49 ± 1.45 ($P=0.017$) from 7.50 ± 2.29 to 6.01 ± 2.88 , -1.44 ± 0.52 ($P=0.000$) from 7.73 ± 3.05 to 6.28 ± 3.42 , -2.18 ± 0.75 ($P=0.000$) from 8.58 ± 3.87 to 6.40 ± 3.48 , and -1.63 ± 1.13 ($P=0.003$) from 8.32 ± 3.35 to 6.69 ± 3.81 respectively (Table 2).

Table 2. Measured myostatin levels (ng/ml) in two studied groups in four different exercises. Asterisk shows statistically significant decrease (P < 0.05)

Group / serum collection		Activity			
Group	Training/Session	Continuous/Water	Discontinuous/Treadmill	Discontinuous/Water	Continuous/Treadmill
Menopause	Pre	9.17±4.38	9.11±4.43	9.69±4.60	9.07±4.14
	Post	7.81±4.21	7.46±4.32	7.61±3.72	7.86±3.53
	t-test	t ₉ = -6.769 P=0.000*	t ₉ = -5.660 P=0.000*	t ₇ = -3.565 P=0.000*	t ₇ = -1.381 P=0.000*
	Covariance	7.32±0.398	7.03±0.398	6.66±0.447	7.47±0.44
Menstrual	Pre	7.50±2.29	7.73±3.05	8.58±3.87	8.32±3.35
	Post	6.01±2.88	6.28±3.42	6.40±3.48	6.69±3.81
	t-test	t ₈ = -3.078 P= 0.017*	t ₈ = -8.374 P= 0.000*	t ₈ = -8.754 P= 0.000*	t ₈ = -4.331 P= 0.003*
	Covariance	7.05±0.442	7.12±0.421	6.48±0.419	6.98±0.419

Table 3. Measured insulin levels (mU/I) in two studied groups in four different activities. Asterisk shows statistically significant decrease (P < 0.05)

Group / serum collection		Activity			
Group	TrainingSession	Continuous/Water	Discontinuous/Treadmill	Discontinuous/Water	Continuous/Treadmill
Menopause	Pre	24.65±18.73	27.54±13.95	26.22±13.91	33.98±13.63
	Post	15.81±11.43	19.89±9.26	15.69±8.51	21.70±11.75
	t-test	t ₉ = -2.671 P= 0.026*	t ₉ = -1.861 P= 0.096	t ₇ = -2.721 P= 0.030*	t ₇ = -4.326 P= 0.003*
	Covariance	17.25±2.69	19.99± 2.68	16.39±3.00	18.79±3.02
Menstrual	Pre	21.48±14.52	24.18±23.22	30.95±18.91	33.82±19.11
	Post	10.37±6.54	14.22±8.49	16.47±16.72	19.98±16.35
	t-test	t ₈ = -3.718 P= 0.006*	t ₈ = -1.743 P= 0.120	t ₈ = -2.931 P= 0.019*	t ₈ = -3.021 P= 0.017*
	Covariance	13.28±2.85	15.87±2.83	14.98±2.83	17.15±2.85

Table 4. Measured glucose levels (mg/dl) in two studied groups in four different exercises. Asterisk shows statistically significant decrease (P < 0.05)

Group / serum collection		Activity			
Group	TrainingSession	Continuous/Water	Discontinuous/Treadmill	Discontinuous/Water	Continuous/Treadmill
Menopause	Pre	80.70±17.29	85.70±16.99	85.87±23.22	84.63±13.14
	Post	66.80±9.34	77.50±16.31	73.00±7.75	72.75±8.08
	t-test	t ₉ = -3.985 P= 0.003*	t ₇ = -1.799 P= 0.115	t ₉ = -2.408 P= 0.039*	t ₇ = -4.395 P= 0.003*
	Covariance	67.34±2.18	76.05±2.19	71.48±2.45	71.72±2.44
Menstrual	Pre	76.55±13.81	77.11±9.70	84.44±16.08	81.88±15.79
	Post	69.88±6.09	66.55±4.36	71.22±9.41	71.66±7.21
	t-test	t ₈ = -1.751 P= 0.118	t ₈ = -3.101 P= 0.015*	t ₈ = -4.726 P= 0.001*	t ₈ = -2.534 P= 0.035*
	Covariance	72.08±2.32	68.53±2.31	70.27±2.30	71.73±2.30

In menopause group, insulin levels in continuous/water, discontinuous/ treadmill, discontinuous/water and continuous/ treadmill groups were decreased -8.84 ± 10.47 (P=0.026)

from 24.65 ± 18.73 to 15.81 ± 11.43 , -7.65 ± 13.00 (P=0.096) from 27.54 ± 13.95 to 19.89 ± 9.26 , -10.53 ± 10.95 (P=0.030) from 26.22 ± 13.91 to

15.69±8.51, and -12.28±8.04 (P=0.003) from 33.98±13.63 to 21.70±11.75 respectively.

In menstrual group, insulin levels in continuous/water, discontinuous/treadmill, discontinuous/water and continuous/treadmill groups were reduced by -11.11±8.96 (P=0.006) from 21.48±14.52 to 10.37±6.54, -9.96±17.16 (P=0.120) from 24.18±23.22 to 14.22±8.49, -14.47±14.81 (P=0.019) from 30.95±18.91 to 16.47±16.72, and -13.84±13.74 (P=0.017) from 33.82±19.11 to 19.98±16.35 respectively (Table 3).

In menopause group, glucose levels in continuous/water, discontinuous/treadmill, discontinuous/water and continuous/treadmill groups were decreased -13.90±11.03 (P=0.003) from 80.70±17.29 to 66.80±9.34, -8.20±10.76 (P=0.115) from 85.70±16.99 to 77.50±16.31, -12.87±20.23 (P=0.039) from 85.87±23.22 to 73.00±7.75, and -11.87±7.64 (P=0.003) from 84.63±13.14 to 72.75±8.08 respectively.

In menstrual group, glucose levels in continuous/water, discontinuous/treadmill, discontinuous/water and continuous/treadmill groups were decreased -6.66±11.42 (P=0.118) from 76.55±13.81 to 69.88±6.09, -10.55±10.21 (P=0.015) from 77.11±9.70 to 66.55±4.36, -13.22±8.39 (P=0.001) from 84.44±16.08 to 71.22±9.41, and -10.22±12.10 (P=0.035) from 81.88±15.79 to 71.66±7.21 respectively (Table 4).

Conclusion:

Obesity is a common phenomenon among menopause women, especially central obesity. Obesity is associated with an increased risk of insulin resistance and type 2 diabetes by increasing concentrations of insulin and glucose (14). Also, myostatin, a negative regulator of muscular growth, is increased in older age and should be taken into account as a potential causative reason for health problems in older men and women. Although the effects of exercise training on myostatin expression in skeletal muscle have been discussed by several studies, the results remain disputable. For example, both increasing and decreasing levels of myostatin mRNA has been seen after different types of exercise training in healthy persons (15). Thus, we decided to design the present study to elucidate the

response of myostatin, insulin and glucose to discontinuous and continuous training on treadmill and in water in two groups of middle aged menopause and menstrual women. Given t-test analysis, the main findings from this investigation were that all myostatin, insulin and glucose reduced after different aerobic training in water and on treadmill.

A) Effects of aerobic exercise on myostatin concentration. With respect to t-test analysis, myostatin levels were significantly decreased after continuous and discontinuous training in water, and discontinuous training on treadmill in menopause group (P<0.05). In contrast, myostatin did not decrease significantly after continuous training on treadmill in menopause group (P>0.05). Meanwhile, myostatin level showed significant decrease in all four different conditions of training in menstrual group (P<0.05).

Given covariance analysis results for myostatin levels in menstrual and menopause groups, the lowest myostatin response was found in discontinuous training in water in both menopause and menstrual groups, while the highest myostatin response belonged to menopause group who conducted continuous training on treadmill. These results were statistically meaningless (P>0.05).

These results are in accordance with previous study which imply on significant reduction of myostatin mRNA contents by training in water for 4 weeks (8). Also, our finding that aerobic exercise is associated with a reduction in serum myostatin levels in women with age of older than 40 years old is consistent with Konopka et al. (9), who recently reported that aerobic exercise training performed for 12 weeks can reduce myostatin mRNA expression in older women (P<0.05).

Although reduction of myostatin levels after different aerobic training protocols varies in menstrual and menopause groups, data from our laboratory suggest that aerobic exercise in general can play a critical role in myostatin reduction and consequently decline the risk of sarcopenia and its associated diseases in older women (15).

B) Effects of aerobic exercise on insulin concentration. Secretion of insulin is mainly controlled by plasma glucose concentration, and the hormone has a number of important metabolic actions. Its main function is to control the uptake

and utilization of glucose in peripheral tissues by the glucose transporter. Insulin concentrations are severely reduced in insulin-dependent diabetes mellitus (IDDM), while its levels are raised in non-insulin-dependent diabetes mellitus (NIDDM) and obesity (16). Serum insulin levels have been used as an estimate of insulin sensitivity (14).

With regard to t-test analysis, our findings in this study present that insulin levels were considerably decline after continuous and discontinuous training in water, and continuous training on treadmill in both menstrual and menopause groups ($P < 0.05$), while no substantial decrease was observed in insulin levels after discontinuous training on treadmill, in spite of a medium decrease ($P > 0.05$).

Considering covariance analysis results for insulin levels in menstrual and menopause groups, the lowest insulin response was detected in continuous and discontinuous training in water in menstrual group, while the highest insulin response belonged to menopause group who performed discontinuous training on treadmill. These results were statistically meaningless ($P > 0.05$).

Overall, increasing oxidative enzymes and glucose transporters in the muscle by exercise training can be considered as an explanation for insulin reduction after aerobic training in both menstrual and menopause overweight women. Our findings are in concordance with Donnelly et al. reported significant decreases in insulin concentrations in both discontinuous and continuous exercise (17).

C) Effects of aerobic exercise on glucose concentration. Glucose is one of the most important carbohydrates in body and considered as the primary source of energy for the body's cells. Glucose levels are tightly regulated in the human body. The determination of glucose levels in blood is critical in the control of diabetes (18).

With respect to t-test analysis, glucose levels were significantly decreased after continuous and discontinuous training in water, and continuous training on treadmill in menopause group ($P < 0.05$). Moreover, glucose levels were declined after discontinuous training on treadmill in menopause group, although this did not reach significance ($P > 0.05$). In menstrual group, glucose levels were significantly decreased after continuous

and discontinuous training on treadmill, and discontinuous training in water ($P < 0.05$). On the contrary, no noticeable drop was detectable after continuous training in water ($P > 0.05$).

Regarding covariance analysis results for glucose levels in menstrual and menopause groups, the lowest glucose response was seen in continuous training in water in menopause group, while the highest glucose response belonged to menopause group who committed discontinuous training on treadmill. These results were statistically meaningless ($P > 0.05$).

These results suggest that aerobic exercise (in water or on treadmill) can reduce the risk of metabolic disease associated with obesity, such as high insulin and glucose concentrations (19).

Evans et.al reported that endurance exercise training was associated with statistically significant improvement in glucose tolerance and reduction in insulin resistance in older adults (20). In comparison, our data show that aerobic training can also reduce insulin and glucose levels in serum. Another study by Ryan et.al describing a significant decrease in plasma glucose and insulin after aerobic and resistive training combined with weight loss in overweight postmenopausal women (21). These data are in support of our findings that demonstrated a drop in insulin and glucose after aerobic training.

It can be concluded that aerobic training provided considerable response in measured factors in studied women groups, in spite of some variations. The lowest myostatin response was seen in discontinuous training in water in both menopause and menstrual groups, while the highest myostatin response observed in menopause group who committed continuous training on treadmill. In summary, this study suggested that discontinuous than continuous exercise, and exercise in the water than on treadmill, Lead to better response in variables are listed.

References:

1. Carr MC. The emergence of the metabolic syndrome with menopause. *J Clin Endocrinol Metab.* 2003;88(6):2404-2411.
2. Toth MJ, Sites CK, Eltabbakh GH, Poehlman, ET. Effect of menopausal status on insulin-stimulated glucose disposal: comparison of middle-aged premenopausal and early postmenopausal women. *Diabetes Care.* 2000;23(6):801-806.
3. Deibert P, König D, Vitolins MZ, Landmann U, Frey Z, Zahradnik HP, et al. Effect of a weight loss intervention on anthropometric measures and metabolic risk factors in pre-versus postmenopausal women. *Nutr J.* 2007;6(31):1-7.
4. Després JP, Nadeau A, Tremblay A, Feriand M, Mooriani S, Lupien PJ, et al. Role of deep abdominal fat in the association between regional adipose tissue distribution and glucose tolerance in obese women. *Diabetes.* 1989;38(3):304-309.
5. Schmitz K, Jensen M, Kugler K, Jeffery R, Leon A. Strength training for obesity prevention in midlife women. *Int J Obes Relate Metab Disord.* 2003;27(3):326-333.
6. Fairey AS, Courneya KS, Field CJ, Bell GJ, Jones LW, Mackey JR. Effects of Exercise Training on Fasting Insulin, Insulin Resistance, Insulin-like Growth Factors, and Insulin-like Growth Factor Binding Proteins in Postmenopausal Breast Cancer Survivors a Randomized Controlled Trial. *Cancer Epidemiol Biomarkers Prev.* 2003;12(8):721-727.
7. Ryan AS, Ivey FM, Prior S, Li G, Hafer-Macko C. Skeletal muscle hypertrophy and muscle myostatin reduction after resistive training in stroke survivors. *Stroke.* 2011;42(2):416-420.
8. Matsakas A, Bozzo C, Cacciani N, Caliare N, Caliaro F, Reggiani C, Mascarello F, et al. Effect of swimming on myostatin expression in white and red gastrocnemius muscle and in cardiac muscle of rats. *Exp Physiol.* 2006;91(6):983-994.
9. Konopka AR, Douglass MD, Kaminsky LA, Jemiolo B, Trappe TA, Trappe S, et al. Molecular adaptations to aerobic exercise training in skeletal muscle of older women. *J Gerontol Biol Sci Med Sci.* 2010;65(11):1201-1207.
10. Sandhu JS, Trivedi SS, Shenoy S. A correlation study between plasma myostatin and peak torque strength gains of bilateral knee extensors and flexors after resistance training in healthy Asian Indians. *J Musculoskelet Res.* 2010;13:109-117.
11. Hukshom CJ, van Dielen FM, Buurman WA, Westerterp-Plantinga MS, Campfield LA, Saris WH. The effect of pegylated recombinant human leptin (PEG-OB) on weight loss and inflammatory status in obese subjects. *Int J Obes Relat Metab Disord.* 2002;26(4):504-509.
12. Nagaraja P, Honnur Krishna H, Shivakumar A, Shrestha AK. Development of quantitative enzymatic method and its validation for the assay of glucose in human serum. *Clin Biochem.* 2012;45(1-2):139-143.
13. Erejuwa OO, Sulaiman SA, Wahab AB, Sirajudeen KN, Salleh MS, Gurtu S. Glibenclamide or metformin combined with honey improves glycemic control in streptozotocin-induced diabetic rats. *Int J Biol Sci.* 2011;7(2):244-252.
14. Frank LL, Sorensen BE, Yasui Y, Tworoger SS, Schwartz RS, Ulrich CM, et al. Effects of exercise on metabolic risk variables in overweight postmenopausal women: a randomized clinical trial. *Obes Res.* 2005;13(3):615-625.
15. Lenk K, Schur R, Linke A, Erbs S, Matsumoto Y, Adams V, et al. Impact of exercise training on myostatin expression in the myocardium and skeletal muscle in a chronic heart failure model. *Eur J Heart Fail.* 2009;11(4):342-348.
16. Rajpathak SN, Gunter MJ, Wylie-Rosett J, Hog Y, Kaplan RC, Muzumdar R, et al. The role of insulin-like growth factor-I and its binding proteins in glucose homeostasis and type 2 diabetes. *Diabetes Metab Res Rev.* 2009;25(1):3-12.
17. Donnelly J, Jacobsen D, Heelan KS, Seip R, Smith S. The effects of 18 months of intermittent vs. continuous exercise on aerobic capacity, body weight and composition, and metabolic fitness in previously sedentary, moderately obese females. *Int J Obes Relat Metab Disord.* 2000;24(5):566-572.
18. Patel A, Macmahon S, Chalmers J, Billot L, Woodward M, Marre M, et al. Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med.* 2008;358(24):2560-2572.
19. Nassis GP, Papantakou K, Skenderi K, Triandafillopoulou M, Kavouras SA, Vannakouilia M, et al. Aerobic exercise training improves insulin sensitivity without changes in body weight, body fat, Adiponectin, and inflammatory markers in overweight and obese girls. *Metabolism.* 2005;54(11):1472-1479.

20. Evans EM, Van Pelt RE, Binder EF, Williams DB, Eshani AA, Kohrt WM. Effects of HRT and exercise training on insulin action, glucose tolerance, and body composition in older women. *J Appl Physiol.* 2001;90(6):2033-2040.
21. Ryan A, Nicklas B, Berman D, Elahi D. Adiponectin levels do not change with moderate dietary induced weight loss and exercise in obese postmenopausal women. *Int J Obes Relat Metab Disord.* 2003;27(9):1066-1071.