

## Research Paper

## The Effect of Varicocelelectomy on Sex Hormone Levels and Semen Parameters



Ali Mohammad Falahati<sup>1</sup>, Mohammad Mohajer-Bastami<sup>1</sup>, Mohammad Natami<sup>1</sup>, Alireza Sobhani<sup>2</sup>, Elahe Taqvaei<sup>1</sup>, Masoomeh Latifi<sup>1</sup>, Soghra Fallahi<sup>\*</sup>

1. Fertility and Infertility Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran.

2. Department of Pathology, Shariati Hospital, Hormozgan University of Medical Sciences, Bandar Abbas, Iran.



**Citation** Falahati AM, Mohajer-Bastami M, Natami M, Sobhani A, Taqvaei E, Latifi M, et al. The Effect of Varicocelelectomy on Sex Hormone Levels and Semen Parameters. *Hormozgan Medical Journal*. 2023; 27(4):187-194. <http://dx.doi.org/10.32598/hmj.27.4.8152>

**doi** <http://dx.doi.org/10.32598/hmj.27.4.8152>



## Article info:

**Received:** 15 Mar 2023

**Accepted:** 29 Jun 2023

**Available Online:** 01 Oct 2023

## Keywords:

Varicocele, Varicocelelectomy, Sexual hormones, Semen

## ABSTRACT

**Objectives:** Varicocelelectomy is a surgical intervention to treat varicocele. It may recover fertility in men by improving semen parameters and sex hormones. This research studied the role of varicocelelectomy, focusing on semen properties and sex hormones.

**Methods:** In this cross-sectional study, the data were collected from 45 patients with different varicocele grades who underwent varicocelelectomy. The data were collected and compared before and after clinical examination.

**Results:** After varicocelelectomy, a significant improvement was observed in certain semen parameters, such as motility and concentration. However, among sex hormones, only testosterone increased after the surgical intervention ( $P=0.007$ ).

**Discussion:** To our knowledge, varicocele treatment may almost improve male fertility. Accordingly, varicocelelectomy positively affects parameters such as testosterone and serum FSH levels and sperm count and motility. As such, varicocelelectomy emerges as a practical approach to improve male fertility and sex disorders.

\* Corresponding Author:

Soghra Fallahi, Assistant Professor.

Address: Fertility and Infertility Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran.

E-mail: [sfallahi@hums.ac.ir](mailto:sfallahi@hums.ac.ir)

## Introduction

**M**ale infertility is a highly prevalent disorder, with several factors involved, the most important of which is varicocele. It can be considered an essential factor in causing the early testicular disorder, marked by an obstruction in the genital tract [1]. Varicocele is a complication caused by the anomalies of veins inside the testicles, resulting in the blockage of blood and subsequent swelling [2]. An overall prevalence of varicocele varies between 15% and 45% in men, with clinical manifestation typically appearing at puberty. The prevalence is estimated to be about 1% before adolescence, increasing to 2% to 16% in adolescence and peaking at 20% by age 15 [3]. Several underlying factors have been reported, including oxidative stress, hypoxia, and hormone imbalance. The primary mechanism responsible for varicocele-induced damage is attributed to DNA injury in the sperm head due to oxidative stress (with high levels of active oxygen species) or reduction of the total antioxidant capacity [4].

Moreover, varicocele progression during adolescence may harm the male reproductive system. Varicocele may increase or decrease in testis size [5]. Once varicocele occurs, prominent features, such as reduced sperm count, decreased testosterone levels, and diminished size (in the same area), are often observed [6]. Varicocele can be associated with a distinct or obscure pain in the testicles or the groin [7]. Other symptoms include a heavy testicular sensation, testicular atrophy, benign prostatic hyperplasia, and other urinary problems [8].

The underlying factors for varicocele are not fully known. Suggested contributors include hyperthermia, testicular blood flow, intravenous pressure, hormonal disorders, renal reflux, autoimmune disorders, and oxidative stress [9]. Hormonal disorders, such as testosterone hormone fluctuations, have been reported in varicocele [10]. Surgical treatment through varicocelectomy can affect serum testosterone levels in the affected cases.

Besides, affected men are at a risk of androgen deficiency [11]. Several studies have shown that varicocelectomy can significantly correct sperm parameters and sex hormones, including the serum testosterone level [12]. Varicocelectomy is a surgical procedure to improve fertility in men. This surgical procedure aims to treat varicocele by preventing the intravenous drainage of the reflux into the testicle and maintaining arterial flow and lymph drainage [13]. Evidence shows that varicocelectomy can improve postoperative testosterone levels,

with a concomitant facilitative (and somewhat satisfactory) impact on sperm parameters [14]. We aimed to investigate the effect of microsurgical varicocelectomy on sex hormones in affected men. This study assessed the role of varicocelectomy, focusing on semen properties and sex hormones.

## Materials and Methods

The present cross-sectional study was conducted in Bandar Abbas City, Iran, from December 17 to March 15, 2019. The study included male patients aged between 22-40 years. A total of 45 patients with varicocele Grade 2 or above, presenting at the Infertility and Urology Center with complaints of infertility or genital pain, underwent examination by a urologist. After diagnosis, patients with a complete medical history, physical examinations, and hormonal profiles were included in the study. After signing an informed consent letter, a demographic profile and clinical data were collected for each patient. The exclusion criteria were affliction with conditions such as testicular atrophy, genitourinary infection, cryptorchidism, testicular trauma, post-genital surgery, hyperprolactinemia, testosterone replacement therapy or antiestrogen, aromatase inhibitors, uncontrolled diabetes, and hypertension. The varicocele grade was determined based on Dubin and Amelar's system.

In grade 1, or low intensity, varicocele is not palpable while examining the testicles but becomes palpable after the Valsalva maneuver. In grade 2 or moderate intensity, varicocele is not visible but is palpable without a Valsalva maneuver, and in grade 3, it is visible through the scrotum without a Valsalva maneuver [15].

Microsurgical varicocelectomy was performed by the method described by Zini et al. [16]. The internal and external spermic vessels were blocked while keeping the elliptic and lymphatic arteries open. Sperm samples were collected through self-stimulation after 3 days of abstinence. Semen analysis assessed parameters such as concentration, viability, motility percentage, sperm shape, and morphology based on World Health Organization (WHO) criteria.

An endocrinologic evaluation was done for serum testosterone levels, FSH, and LH evaluation, followed by electrochemiluminescence. The spermogram data and preoperative sexual hormone values were compared with the data 6 months after varicocelectomy. The data were analyzed in SPSS software, version 16 through paired-sample t-test and Wilcoxon test.  $P < 0.05$  were considered the significant data interpretation level.

## Results

In this study, 45 male participants with varicocele were included. Their mean age was  $29.5 \pm 5.7$  years. The demographic data showed that 26.7% of patients had a smoking history, and 22.2% reported consanguinity with their spouses (Table 1).

The results showed that most participants (86.7%) had varicocele grade 2, with only 13.3% presenting with varicocele Grade 3 (Table 2).

According to the results, the mean follicle stimulating hormone (FSH) level before and after the intervention was  $9.83 \pm 11.75$  IU/L and  $8.72 \pm 10.04$  IU/L, respectively (Table 3). The Wilcoxon test results showed no statistically significant difference between these two phases ( $P=0.069$ ). Similarly, the mean luteinizing hormone (LH) levels before and after the intervention were  $6.55 \pm 4.25$  and  $7.07 \pm 6.20$  IU/L, respectively, with no statistically significant difference observed ( $P=0.820$ ). Additionally, the mean prolactin levels before and after the intervention were  $373.70 \pm 180.08$  and  $342.43 \pm 284.25$

$\mu\text{IU/mL}$ , respectively, without statistically significant difference ( $P=0.670$ ).

The results showed that the mean testosterone levels were  $3.02 \pm 1.33$  and  $4.06 \pm 1.51$  ng/mL before and after the intervention, respectively, presenting a statistically significant difference ( $P=0.007$ ).

The results showed that the mean volumes of semen, before and after the intervention, were  $2.32 \pm 1.32$  and  $3.15 \pm 1.22$  mL, respectively. The difference between the pre-test and post-test was not statistically significant ( $P=0.054$ ). Also, the pH value remained constant at 7.8 before and after the intervention.

Based on the results, the mean progressive motility was  $33.87\% \pm 23.34\%$  before the intervention and  $40.95\% \pm 21.31\%$  after the intervention, with no statistically significant difference between these two phases ( $P=0.136$ ). According to the findings, the mean sperm motility index was  $108.53 \pm 104.66$  before the intervention, changing to  $104.40 \pm 72.56$  after the intervention. The difference between the pre- and post-test was not statistically significant ( $P=0.053$ ).

**Table 1.** Distribution of demographic data

Personal Data		Mean $\pm$ SD/No. (%)
Age (y)		29.5 $\pm$ 5.7
Consanguinity	Yes	10(22.2)
	No	35(77.8)
Family history of infertility	Yes	6(13.3)
	No	39(86.7)
Smoking	Yes	12(26.7)
	No	33(73.3)
Addiction	Yes	6(13.3)
	No	39(86.7)

**Table 2.** Distribution of varicocele status

Varicose Status		No. (%)
Grade	2	39(86.7)
	3	6(13.3)
Position	Left	39(86.7)
	Bilateral	6(13.3)

**Table 3.** Sex hormone profile before and after intervention

Hormones	Mean±SD			P
	Before	After	Difference	
FSH (IU/L)	9.83±11.75	8.72±10.04	1.11±3.99	0.069
LH (IU/L)	6.55±4.25	7.07±6.20	0.52±3.71	0.820
Prolactin (μIU/mL)	373.70±180.08	342.43±284.25	31.27±278.47	0.670
Testosterone (ng/mL)	3.02±1.33	4.06±1.51	-1.04±1.29	0.007*

FSH: Follicle stimulating hormone; LH: Luteinizing hormone.

\*P<0.05.

Furthermore, the mean total sperm motility was  $58.77 \pm 70.13$  million before the intervention. However, it increased to  $64.68 \pm 74.08$  after the intervention ( $P=0.001$ ). The results also showed that the mean total functional sperm concentrations before and after the intervention were  $40.09 \pm 50.39$  and  $42.06 \pm 51.26$  million, respectively (Table 4), showing a statistically significant difference ( $P=0.003$ ).

## Discussion

Varicocele can adversely affect spermatogenesis by increasing testicular temperature and refluxing toxic metabolites [17]. Varicolectomy is a surgical inter-

vention for treating male infertility in male adults and adolescents. Moreover, it treats testicular hemorrhage and hypogonadism and improves sex hormones such as testosterone [18]. The present study demonstrated that varicolectomy may have therapeutic effects on infertile patients. Our finding showed that testosterone level was 3.01 ng/mL before surgery, but after surgery, a significant elevation of up to 4.06 ng/mL was observed ( $P=0.007$ ). Several studies reported that men with total testosterone below 400 ng/dL (4 ng/mL) exhibited higher serum testosterone levels than those with normal total testosterone [19]. However, other studies showed no significant increase in testosterone levels after varicolectomy [20].

**Table 4.** Semen parameters before and after the intervention

Variables	Mean±SD			P
	Before	After	Difference	
Semen volume (mL)	2.32±1.32	3.15±1.22	0.83±1.53	0.054
pH	7.80±0.00	7.80±0.00	0.00±0.00	---
Total alive concentration ( $10^6$ /mL)	39.39±43.67	33.54±29.65	5.85±43.86	0.613
Progressive motility (%)	33.87±23.34	40.95±21.31	-7.08±17.32	0.136
Normal morphology (%)	26.27±13.92	30.07±12.98	-3.80±11.78	0.232
Motile sperm concentration ( $10^6$ /mL)	21.93±31.01	24.00±31.49	-2.07±1.48	0.001*
Functional sperm concentration ( $10^6$ /mL)	18.72±24.07	20.55±25.55	-1.83±1.65	0.001*
Sperm motility index	108.53±104.66	104.40±72.56	4.13±57.20	0.053
All sperm (million)	100.97±117.05	110.38±121.93	-9.41±13.47	0.001*
Total motile sperm (million)	58.77±70.13	64.68±74.08	-5.91±5.55	0.001*
Total functional sperm concentration (million)	40.09±50.39	42.06±51.26	-1.97±2.08	0.003*

\*P<0.05.

In addition, similar results have been obtained in other studies [21, 22]. Resorlu et al. did not report any change in serum total testosterone levels in infertile patients after surgery. This finding was consistent with the study by Such et al. Consequently, this surgery may decrease the odds of abnormalities in the Leydig cells. These researchers showed that age is not a significant contributor to varicocele surgery and the high success rate across different ages [23]. However, varicocele surgery in older patients with hypogonadism has been ended with more satisfactory results [19].

In another study by Ishikawa et al., it was noted that the serum testosterone levels are insufficient to estimate elliptic androgens' production. While the serum testosterone level was normal in patients with varicocele, the production of elliptic testosterone decreased in patients with varicocele. In this study, the increased serum-free testosterone levels after varicocele surgery could increase sperm concentration and motility [24]. There is no significant relationship between the mechanism showing how varicocele affects testosterone synthesis. However, the mechanisms that might underlie the reduced activity of 17 and 20 desmolase and 17  $\alpha$ -hydroxylase and disrupted enzymes could increase the testicular temperature and disorder the Leydig cell response to defective gonadotropin stimulation [25].

Also, the varicocele surgery and its effect on sperm parameters were evaluated in this study. Three studies identified a significant correlation between the correction of some sperm parameters after varicocele surgery. For instance, in the total motility parameter, the amount of movement before the intervention was 58.77 million, and after the intervention, 64.68 million ( $P=0.001$ ). The total functional sperm concentration ( $P=0.003$ ) was 40.09 before the intervention and increased to 42.06 after the intervention. In this study, the total concentration of sperm movement before was 21.93 and reached 24.00 after the intervention. The difference was statistically significant. Varicocele surgery can play a corrective role in the performance of sperm parameters. Several studies showed that sperm motility can be significantly increased compared to preoperative surgery. At the same time, the correction of morphology and count did not reveal any statistically significant differences [26].

In another study by Chu et al., varicocele surgery affected several sperm parameters. In this study, the total motile count was significantly increased and improved. The count was 2.8 million before the surgery, reaching 18.2 million after the varicocele surgery [27]. Shabana et al. also reported that varicocele surgery could significant-

ly change and increase the density and progression of sperm motility [28].

In contrast, the FSH and LH hormones did not change significantly, and the observed alterations in these hormones were minimal. Furthermore, the upward trend of testosterone was also effectively increased [28]. Other studies have shown that FSH and LH were significantly reduced in patients with hypogonadism, which may be due to an increase in total testosterone levels due to the improved function of Leydig cells in the follow-up. Previous studies did not show a specific effect of varicocele surgery on serum LH, even when the mean serum total testosterone improved [21]. Leydig cell disorder leads to an increase in FSH and LH in infertile men with varicocele. Fujisawa et al. (2006) maintained that patients undergo varicocele surgery with decreased FSH and LH levels [29].

In our study, the level of FSH and LH before and after varicocele surgery also changed. However, the difference was not statistically significant, and the level of FSH before and after surgery was 9.83 and 8.72 IU/L, respectively, similar to other studies. However, the LH levels before and after the surgery were 6.55 and 7.07 IU/L, respectively, which is not consistent with previous studies. Chen et al. showed that the essential factor to ascertain the success of varicocele surgery is the low serum FSH concentration [30].

On the contrary, Chen et al. showed no significant difference in FSH levels and fertility rates after varicocele surgery [30]. Fuping Li reported that the mean serum testosterone level increased after varicocele surgery, accompanied by a decrease in the level of FSH and LH after surgery. This pattern aligns with the negative feedback mechanism of the hypothalamic-pituitary-gland axis [31]. Kaneko et al. showed that after varicocele surgery, sperm volume increased in adolescents and adults. In patients with improved sperm volume, FSH level was significantly reduced. The serum FSH levels may play a predictive role in post-surgery testicular function [32]. Su et al. reported that although varicocele surgery improved serum testosterone, no difference was seen between serum FSH and LH levels or the relationship between varicocele grade and serum testosterone levels. Also, in terms of semen parameters, sperm motility was significantly increased compared to the sperm count after surgery, which can be explained by the recovery of testosterone synthesis needed for epididymis function [33].

In another study, Fujisawa showed that the quality of sperm increased in varicocele patients after varicocelectomy, but no significant difference was found in testosterone levels [34]. In a study by Plymate et al. on patients with varicocele, a significant increase was seen in serum FSH before surgery, and the quality of semen was impaired, too. When the serum FSH levels decreased, the function of Sertoli cells could be improved, which results from surgery [35].

A proposed hypothesis regarding the effective role of varicocelectomy suggests a post-surgical improvement of testicular environmental temperature [36]. Increasing the testicular scrotum temperature decreases the concentration of 5 $\alpha$ reductase, associated with converting testosterone to 5 $\alpha$ -dihydrotestosterone, potentially causing complications for varicocele in patients. Varicocelectomy corrects this problem [37]. The Leydig cell function is to produce testosterone under the control of LH. FSH initially develops testosterone production. This process protects LH, and this function requires the regular operation of the hypothalamic-pituitary-gland axis [38].

In our study, it has been shown that varicocelectomy, as in most previous studies, has a statistically significant positive effect on testosterone. Moreover, our study results were consistent with most other results regarding the FSH hormone, although this observation was not significant. In the sperm parameter analysis, our study showed a significant positive effect on sperm motility after the surgery.

It should be noted that the present research faced several limitations, such as the lack of cooperation by some patients, the need to substitute them with other cooperative patients, and the time-consuming and exhaustive duration of research. To obtain more comprehensive information, it is recommended to conduct cohort studies.

## Conclusion

In light of the present findings, varicocelectomy positively affected parameters such as serum testosterone, FSH levels, sperm count, and motility. Varicocelectomy is an effective intervention to improve fertility and sexual dysfunction in infertile men.

## Ethical Considerations

### Compliance with ethical guidelines

The present study was approved by the Ethics and Human Rights Committee of Hormozgan University of Medical Sciences (Code: IR.HUMS.REC.1394.193).

## Funding

The research was funded by the Student Research Committee, Hormozgan University of Medical Science.

## Authors' contributions

Study design and administration: Soghra Fallahi; Data collection, manuscript draft preparation: Ali Mohammad Falahati; Data collection: Mohammad Mohajer-Bastami and Elahe Taqvae; Clinical examination: Mohammad Natami; Experiments: Alireza Sobhani; Resources, review and editing: Masoomeh Latifi.

## Conflict of interest

The authors declared no conflict of interest.

## Acknowledgments

The researchers thank all those who participated in this research, especially those working in the Special Bandar Abbas Clinic. We also appreciate Dr. Sobhani's Medical Diagnostic Laboratory.

## References

- [1] Kantartzi P, Goulis CD, Goulis G, Papadimas I. Male infertility and varicocele: myths and reality. *Hippokratia*. 2007; 11(3):99-104. [PMID]
- [2] Evers J, Collins J, Clarke J. Surgery or embolisation for varicoceles in subfertile men. *Cochrane Database Syst Rev*. 2009; 1. [DOI:10.1002/14651858.CD000479.pub4]
- [3] Vujkovic M, de Vries JH, Dohle GR, Bonsel GJ, Lindemans J, Macklon NS, et al. Associations between dietary patterns and semen quality in men undergoing IVF/ICSI treatment. *Hum Reprod*. 2009; 24(6):1304-12. [DOI:10.1093/humrep/dep024] [PMID]
- [4] Choi WS, Kim SW. Current issues in varicocele management: A review. *World J Mens Health*. 2013; 31(1):12-20. [Link]
- [5] Sayfan J, Siplovich L, Koltun L, Benyamin N. Varicocele treatment in pubertal boys prevents testicular growth arrest. *J Urol*. 1997; 157(4):1456-7. [DOI:10.1016/S0022-5347(01)65021-3] [PMID]
- [6] Said S, Aribarg A, Virutamsen P, Chutivongse S, Koetsawang S, Meherjee P, et al. The influence of varicocele on parameters of fertility in a large group of men presenting to infertility clinics. *Int J Obstet Gynecol*. 1993; 40(3):274. [DOI:10.1016/0020-7292(93)90883-X]
- [7] Masson P, Brannigan RE. The varicocele. *Urol Clin North Am*. 2014; 41(1):129-44. [DOI:10.1016/j.ucl.2013.08.001] [PMID]



- [8] Gat Y, Gornish M, Heiblum M, Joshua S. Reversal of benign prostate hyperplasia by selective occlusion of impaired venous drainage in the male reproductive system: novel mechanism, new treatment. *Andrologia*. 2008; 40(5):273-81. [DOI:10.1111/j.1439-0272.2008.00883.x] [PMID]
- [9] Naughton CK, Nangia AK, Agarwal A. Pathophysiology of varicoceles in male infertility. *Hum Reprod Update*. 2001; 7(5):473-81. [DOI:10.1093/humupd/7.5.473] [PMID]
- [10] Damsgaard J, Joensen UN, Carlsen E, Erenpreiss J, Blomberg Jensen M, Matulevicius V, et al. Varicocele is associated with impaired semen quality and reproductive hormone levels: A study of 7035 healthy young men from six European Countries. *Eur Urol*. 2016; 70(6):1019-29. [DOI:10.1016/j.eururo.2016.06.044] [PMID]
- [11] Tanrikut C, Goldstein M, Rosoff JS, Lee RK, Nelson CJ, Mulhall JP. Varicocele as a risk factor for androgen deficiency and the effect of repair. *BJU Int*. 2011; 108(9):1480-4. [DOI:10.1111/j.1464-410X.2010.10030.x] [PMID]
- [12] Almahdy AEM, Eldin AAG, Abdullah MM, Abuzaid MI. Varicocele repair outcome with respect to hormonal profile and spermogram pattern. *Menoufia Med J*. 2014; 27:164-8. [DOI:10.4103/1110-2098.132792]
- [13] Binsaleh S, Lo KC. Varicocelectomy: Microsurgical inguinal varicocelectomy is the treatment of choice. *Can Urol Assoc J*. 2007; 1(3):277-8. [PMID]
- [14] Hsiao W, Rosoff JS, Pale JR, Powell JL, Goldstein M. Varicocelectomy is associated with increases in serum testosterone independent of clinical grade. *Urology*. 2013; 81(6):1213-8. [DOI:10.1016/j.urology.2013.01.060] [PMID]
- [15] Dubin L, Amelia RD. Etiologic factors in 1294 consecutive cases of male infertility. *Fertil Steril*. 1971; 22(8):469-74. [DOI:10.1016/S0015-0282(16)38400-X] [PMID]
- [16] Zini A, Fischer A, Bellack D, Noss M, Kamal K, Chow V, et al. Technical modification of microsurgical varicocelectomy can reduce operating time. *Urology*. 2006; 67(4):803-6. [DOI:10.1016/j.urology.2005.10.044] [PMID]
- [17] Kantartzi PD, Goulis ChD, Goulis GD, Papadimas I. Male infertility and varicocele: Myths and reality. *Hippokratia*. 2007; 11(3):99-104. [PMID]
- [18] Mehta A, Goldstein M. Microsurgical varicocelectomy: A review. *Asian J Androl*. 2013; 15(1):56-60. [DOI:10.1038/aja.2012.98] [PMID] [PMCID]
- [19] Hsiao W, Rosoff JS, Pale JR, Greenwood EA, Goldstein M. Older age is associated with similar improvements in semen parameters and testosterone after subinguinal microsurgical varicocelectomy. *J Urol*. 2011; 185(2):620-5. [DOI:10.1016/j.juro.2010.09.114] [PMID]
- [20] Rodriguez Peña M, Alescio L, Russell A, Lourenco da Cunha J, Alzu G, Bardoneschi E. Predictors of improved seminal parameters and fertility after varicocele repair in young adults. *Andrologia*. 2009; 41(5):277-81. [DOI:10.1111/j.1439-0272.2009.00919.x] [PMID]
- [21] Ozden C, Ozdal OL, Bulut S, Guzel O, Koyuncu HH, Memis A. Effect of varicocelectomy on serum inhibin B levels in infertile patients with varicocele. *Scand J Urol Nephrol*. 2008; 42(5):441-3. [DOI:10.1080/00365590802028141] [PMID]
- [22] Zheng YQ, Gao X, Li ZJ, Yu YL, Zhang ZG, Li W. Efficacy of bilateral and left varicocelectomy in infertile men with left clinical and right subclinical varicoceles: A comparative study. *Urology*. 2009; 73(6):1236-40. [DOI:10.1016/j.urology.2008.11.050] [PMID]
- [23] Reşorlu B, Kara C, Şahin E, Ünsal A. The significance of age on success of surgery for patients with varicocele. *Int Urol Nephrol*. 2010; 42(2):351-6. [DOI:10.1007/s11255-009-9589-y] [PMID]
- [24] Ishikawa T, Fujisawa M. Varicocele ligation on free testosterone levels in infertile men with varicocele. *Arch Androl*. 2004; 50(6):443-8. [DOI:10.1080/01485010490485803] [PMID]
- [25] Fujisawa M, Hayashi A, Imanishi O, Tanaka H, Okada H, Matsumoto O, et al. The significance of gonadotropin-releasing hormone test for predicting fertility after varicocelectomy. *Fertil Steril*. 1994; 61(4):779-82. [DOI:10.1016/S0015-0282(16)56662-X] [PMID]
- [26] Cakiroglu B, Sinanoglu O, Gozukucuk R. The effect of varicocelectomy on sperm parameters in subfertile men with clinical varicoceles who have asthenozoospermia or teratozoospermia with normal sperm density. *ISRN Urol*. 2013; 2013:698351. [DOI:10.1155/2013/698351] [PMID] [PMCID]
- [27] Chu DI, Zderic SA, Shukla AR, Srinivasan AK, Tasian GE, Weiss DA, et al. Does varicocelectomy improve semen analysis outcomes in adolescents without testicular asymmetry? *J Pediatr Urol*. 2017; 13(1):76. e1-. e5. [DOI:10.1016/j.jpuro.2016.09.010] [PMID] [PMCID]
- [28] Shabana W, Teleb M, Dawod T, Elsayed E, Desoky E, Shaha A, et al. Predictors of improvement in semen parameters after varicocelectomy for male subfertility: A prospective study. *Can Urol Assoc J*. 2015; 9(9-10):E579-82. [DOI:10.5489/cuaj.2808] [PMID] [PMCID]
- [29] Fujisawa M, Dobashi M, Yamasaki T, Kanzaki M, Okada H, Arakawa S, et al. Significance of serum inhibin B concentration for evaluating improvement in spermatogenesis after varicocelectomy. *Hum Reprod*. 2001; 16(9):1945-9. [DOI:10.1093/humrep/16.9.1945] [PMID]
- [30] Chen SS, Chen LK. Predictive factors of successful varicocelectomy in infertile patients. *Urol Int*. 2011; 86(3):320-4. [DOI:10.1159/000322825] [PMID]
- [31] Li F, Yue H, Yamaguchi K, Okada K, Matsushita K, Ando M, et al. Effect of surgical repair on testosterone production in infertile men with varicocele: A meta-analysis. *Int J Urol*. 2012; 19(2):149-54. [DOI:10.1111/j.1442-2042.2011.02890.x] [PMID]
- [32] Kaneko T, Sasaki S, Yanai Y, Umemoto Y, Kohri K. Effect of microsurgical repair of the varicocele on testicular function in adolescence and adulthood. *Int J Urol*. 2007; 14(12):1080-3. [DOI:10.1111/j.1442-2042.2007.01894.x] [PMID]
- [33] Su LM, Goldstein M, Schlegel PN. The effect of varicocelectomy on serum testosterone levels in infertile men with varicoceles. *J Urol*. 1995; 154(5):1752-5. [DOI:10.1016/S0022-5347(01)66776-4]
- [34] Ok F, Erdogan O, Durmus E. Can preoperative gonadotropin and testosterone levels predict the success of varicocelectomy? *Andrologia*. 2020; 52(11):e13887. [PMID]

- [35] Plymate SR, Paulsen CA, McLachlan RI. Relationship of serum inhibin levels to serum follicle-stimulating hormone and sperm production in normal men and men with varicoceles. *J Clin Endocrinol Metab.* 1992; 74(4):859-64. [DOI:10.1210/jcem.74.4.1548351] [PMID]
- [36] Lü YQ, Chen B. [Progress in researches on the mechanism of varicocele-induced male infertility (Chinese)]. *Zhonghua Nan Ke Xue.* 2008; 14(5):454-8. [PMID]
- [37] Aquila S, Montanaro D, Guido C, Santoro M, Perrotta I, Gervasi S, et al. Human sperm molecular anatomy: The enzyme 5 $\alpha$ -reductase (SRD5A) is present in the sperm and may be involved in the varicocele-related infertility. *Histochem Cell Biol.* 2015; 144(1):67-76. [DOI:10.1007/s00418-015-1320-8] [PMID]
- [38] Tian D, Huang W, Yan H, Zong H, Zhang Y. Effect of varicocelectomy on serum FSH and LH levels for patients with varicocele: A systematic review and meta-analysis. *Indian J Surg.* 2018; 80(3):233-8. [DOI:10.1007/s12262-016-1571-1] [PMID] [PMCID]