

## ⇒ Research Article



# Identification of Effective Factors on the Survival of Hemodialysis Patients Using the LASSO-Cox Approach

Shideh Rafati<sup>1</sup>, Laleh Hassani<sup>2\*</sup>, Mohamad Tamaddondar<sup>3</sup>, Abdolnabi Sabili<sup>4</sup><sup>1</sup>Department of Biostatistics and Epidemiology, Kerman University of Medical Sciences, Kerman, Iran<sup>2</sup>Mother and Child Welfare Research Center, Hormozgan University of Medical Sciences, Bandar Abbas, Iran<sup>3</sup>Hormozgan University of Medical Sciences, Bandar Abbas, Iran<sup>4</sup>Education Department, Social Security Organization, Bandar Abbas, Iran

## Abstract

**Background:** Hemodialysis is a method for removing excess fluid and uremic waste from the body of chronic renal failure (CRF) patients. It stops death in patients with CRF. Various factors can affect the survival of patients undergoing hemodialysis treatment. The current study aimed to determine some factors affecting hemodialysis patients' survival using the LASSO-Cox approach.

**Methods:** This cross-sectional study examined 252 patients undergoing hemodialysis from 2010 to 2016 in all hospitals of Bandar Abbas, Iran, whose data had been recorded in the hemodialysis unit, or it was possible to obtain their files. The Cox model under the LASSO variable selection technique (LASSO-Cox) was applied for data analysis. The collected data were analyzed using SPSS 23.0 (IBM) software and glmnet and survival packages in R software, version 3.5.1.

**Results:** Overall, 35 (13.9%) death events were observed in this study. The mean follow-up time of 252 patients was equal to  $10.93 \pm 7.82$  years. Based on the results of the Cox-Lasso method, the risk of mortality for patients with a diploma was 49% lower than that of the illiterate group (HR=0.51; 95% CI: 0.02, 0.57;  $P=0.021$ ). The risk of death for unemployed patients and farmers was 0.66 (HR=1.66; 95% CI: 1.51, 25.79;  $P=0.004$ ) and 0.29 (HR=1.29; 95% CI: 1.14, 27.40;  $P=0.005$ ) higher than that of employees, respectively. Further, by an increase in the frequency of dialysis per week, the risk of death was reduced by 0.34 (HR=0.66; 95% CI: 0.04, 0.82;  $P=0.022$ ), and one unit increase in the duration of dialysis per hour increased the risk of death by 2.23 times (HR=2.23; 95% CI: 1.41, 27.1;  $P=0.010$ ).

**Conclusion:** Education level, job, frequency of hemodialysis per week, and duration of hemodialysis per hour were the most important variables in the survival of hemodialysis patients. As a result, it seems that more education for nurses working in hemodialysis wards and the involved patients can be useful to heed the recommendations of physicians in terms of the required duration for hemodialysis per hour and the number of sessions per week.

**Keywords:** Chronic Kidney Failure, Dialysis, Survival Analysis

## \*Correspondence to

Laleh Hassani,  
Email: 7hassani1969@gmail.com



Received September 30, 2021, Accepted: February 20, 2022, Published Online: November 2, 2022

## Background

An inappropriate and unhealthy lifestyle is responsible for the occurrence of many types of chronic diseases. Chronic illness is the cause of much mortality throughout the world (1). Chronic renal failure (CRF) is one of the chronic diseases that has risen in recent years and has increased tenfold in the United States in the past 20 years (2). According to the statistics of the Center for the Management of Transplantation and Special Diseases of the Ministry of Health of Iran, the annual growth rate of the disease is nearly 11% in this country (3). Patients with CRF are unable to survive without renal replacement therapy (4). According to a report from the above-mentioned center, the number of patients with kidney failure is equal to 320 000 people, 49%, 48%, and 3% of whom use transplantation therapy, hemodialysis, and peritoneal dialysis, respectively (5). Although hemodialysis prevents death in patients with CRF, it does

not treat renal disease and does not compensate for the endocrine and metabolic activities of the kidneys, and the patient is exposed to some problems and complications (6, 7). In the US, diabetes mellitus accounts for more than 45% of newly diagnosed cases with end-stage renal disease (ESRD), and it is also the second most common cause of hypertension, which is estimated to account for 28% of ESRD cases. In addition, the results of a study in Iran indicated that hypertension has been reported as the most common cause of ESRD (7, 8). Serum albumin is one of the ways to assess the status of patients with ESRD, and checking changes in serum levels of albumin can help diagnose the progression of the disease and use appropriate treatment to prevent future risks (8, 9). In studies with large sample sizes, it has been shown that a low concentration of albumin is a reliable predictor of mortality in ESRD patients (10, 11). The inadequacy of hemodialysis is one of the main causes of mortality

in patients with ESRD (12). Several factors such as diet, type of filter, device rotation, hemodialysis time, and underlying disease affect hemodialysis adequacy (13). In hemodialysis, uremic removal is calculated by a number used to quantify the hemodialysis and peritoneal dialysis treatment adequacy (Kt/V) formula with a normal level of 1.2. For this purpose, blood urine nitrogen is controlled in the patient to measure urea removal before and after hemodialysis. Therefore, the amount of urea removal in a 4-hour phase of hemodialysis is important in assessing adequacy, and it does not depend on the plasmatic urea level alone (14). One of the important issues in effective hemodialysis is hemodialysis duration. Based on the findings of a study conducted in this regard, mortality in hemodialysis patients had a direct relationship with hemodialysis duration. The mortality rate of hemodialysis patients who underwent hemodialysis below 10.5 hours of hemodialysis in the first week was far more than that of those who underwent hemodialysis every week for more than 12 hours (15).

Survival analysis is a branch of statistics for data analysis where the outcome variable of interest is the time of occurrence of an event. Due to the importance of identifying factors affecting the survival of dialysis patients, the researchers were encouraged to conduct this study.

### Objectives

The current study sought to determine some factors affecting the lifespan of hemodialysis patients in Bandar Abbas, Iran using survival analysis.

### Methods

In this cross-sectional study, patients' information was collected through a census from the Hemodialysis Department of Shahid Mohammadi, Kodakan, and Khalij-e-Fars Hospitals in Bandar Abbas, Iran. According to studies on the total number of patients who were admitted during 2010-2016, 252 cases were examined, whose data were recorded in the hemodialysis department, and their medical records were available, among whom, only 35 people died by the end of 2016. In the current study, death was considered as the desired event, and excluded cases included live subjects at the end of the study, people missed the follow-up, and kidney transplant recipients. The survival time of patients was calculated from the beginning of hemodialysis to the end of the study in 2016. Data were collected based on a designed checklist which was approved by a specialist. The checklist included variables such as age, gender (female or male), educational status (illiteracy, low literacy, diploma, university graduate, and the number of years of studying), marital status (single, married, or widow), and employment status (housewife, unemployed, civil servant, farmer, retired, or others). The other variables

were smoking status (smoker or non-smoker), the type of underlying disease that leads to hemodialysis (diabetes, hypertension, stones and kidney obstruction, polycystic kidney disease, congenital, unspecified diseases, or others), and age of underlying disease diagnosis. The remaining factors included cardiovascular diseases, family support, age of hemodialysis initiation, mean serum albumin, and serum creatinine. The obtained data were coded and computerized for analysis.

In the present study, excluded cases were live subjects at the end of the study, patients who missed the follow-up, and kidney transplant recipients. The survival time of the patients was calculated by the years from the onset of hemodialysis to the end of the study in 2016. Due to the presence of sparse data (35 cases (13.9%) died and 217 (86.1%) cases were excluded), the performance of ordinary Cox regression is questionable. In such a situation, Cox regression by the Least Absolute Shrinkage and Selection Operator (LASSO) method has been introduced as a suitable approach to finding variables affecting patients' survival.

The collected data were analyzed using SPSS 23.0 (IBM) and installing glmnet and survival packages in R software, version 3.5.1. A two-sided  $P < 0.05$  was considered statistically significant. It is worth noting that in the analysis of survival, conclusions are based on the hazard ratio (HR). If its value is greater than one, it indicates that the group is at higher risk for the occurrence of the event compared to the base group. If its value is less than one, compared to the base group, there will be a lower risk for the occurrence of the desired event, and HR-1 is usually used to express such situations.

### Results

In the current study, 252 hemodialysis patients were investigated who were referred to Shahid Mohammadi, Kodakan, and Khalij-e-Fars hospitals in Bandar Abbas. Among 252 patients, there were 35 (13.9%) and 217 (86.1%) cases of death and exclusion, respectively. The mean of follow-up time of 252 patients was equal to  $10.93 \pm 7.82$  years, and their mean age was  $53.39 \pm 18.09$  years in death or censoring time. Further, the mean age of hemodialysis initiation in patients was equal to  $42.88 \pm 17.07$  years, and the mean score of their body mass index (BMI) was  $22.87 \pm 4.24$ . Most women were housewives (87.1%), and most men were unemployed or retired. No cases with HIV infection were observed in these patients. Hemodialysis lasted 4 hours each time for 194 patients (77%). Overall, 171 (67.9%) patients were hemodialyzed three times a week (Table 1).

The 10-year survival rate of hemodialysis patients was equal to 0.87, implying that 0.87 of the patients will survive more than ten years (Figure 1). The results of the LASSO-Cox model (Table 2) demonstrated that by adjusting other variables, for one unit of increase in BMI, the risk

**Table 1.** Hemodialysis Patients' Characteristics

Variables	Mean	SD	
Follow-up time (y)	10.93	7.82	
Body mass index (kg/m <sup>2</sup> )	22.87	4.24	
Age of dialysis initiation (y)	42.88	17.07	
Age in death or censoring time (y)	53.39	18.09	
	Frequency	Percent	
Blood group	O	108	42.9
	A	62	24.6
	B	72	28.6
	AB	10	3.9
Education level	Illiterate	87	34.5
	Low literacy	118	46.8
	Diploma	40	15.9
	Academic	7	2.8
Job	Unemployed	46	18.2
	Farmer	44	17.5
	Housewife	101	40.1
	Employed	61	24.2
Gender	Men	136	54
	Women	116	46
Marital	Married	207	82.1
Smoking	Yes	91	36.1
Diabetes	Yes	134	53.2
Hypertension	Yes	152	60.3
Urinary stone	Yes	23	9.1
Kidney cyst	Yes	11	4.4
Cardiac pulmonary disease	Yes	50	19.8
Congenital disease	Yes	4	1.6
Glomerulonephritis	Yes	18	7.1
CRF history in family	Yes	24	9.5
Anemia history	Yes	195	77.4
Take of EPREX	Yes	239	94.8
HCV	Yes	8	3.2
HBV	Yes	3	1.2
Kidney transplantation	Yes	27	10.7
Stopped dialyze in reason of kidney function	Yes	22	8.7

Note. SD: Standard deviation; CRF: Chronic renal failure; HCV: Hepatitis C virus; HBV: Hepatitis B virus.

of mortality is reduced by 0.10. The risk of death for those with a diploma was 0.49 lower than that of the illiterate group. Additionally, the risk of death for unemployed patients and farmers was 0.66 and 0.29 higher than that of employers, respectively. Another significant factor in this study was the duration of each hemodialysis session. The risk of death has fallen by 0.34 for one unit of increase in the number of hemodialyses. Moreover, by controlling other factors, increasing the duration of each hemodialysis session increased the hazard of death 2.23 times.

## Discussion

Our study attempted to determine factors affecting the hemodialysis patients' survival using the LASSO-Cox approach. Based on the results of this study, the level of education, occupation, the hemodialysis center, the duration of hemodialysis regarding hours per session, and the frequency of hemodialysis per week were the most important prognostic factors.

The results also represented that by controlling other variables, the risk of death in people with kidney failure was reduced by 10% with one unit increase in the BMI. Weight loss in patients with hemodialysis is the most important goal in every hemodialysis session, and patients are weighed at the beginning and the end of hemodialysis, thus the result of this study suggests a paradox, which has been addressed in previous research (16). It should be noted that BMI reduction is not addressed in patients for two reasons; malnutrition in hemodialysis patients is a major risk in the mortality of patients (17, 18) and depends on receiving insufficient energy. Getting 35 kcal per kg of body weight as the recommended calorie intake will help maintain protein and thus prevent a negative nitrogen balance (19, 20).

Furthermore, the risk of mortality for those with a diploma was 0.49 lower than that of the illiterate group. It may be justified that treatment adherence is higher in these individuals because of higher education or they are more likely to seek information because of their education and even have a better quality of life (21). Literacy by making basic changes in the knowledge of people and attitudes affected health and disease and the other aspects of life, and people with a high level of education have better availability to supportive sources and better control the side effects of diseases, and ultimately have a better quality of life (22). Although it has been proven in some studies, there have been contradictions in this regard (23).

The risk of mortality for unemployed patients and farmers was 0.66 and 0.29 higher than that of employers,




Figure 1. The Kaplan Meier Survival Curve of Patients.

**Table 2.** The Most Important Variables Selected Based on the LASSO-Cox Model

Variables	Coefficient	P Value	95% CI	Hazard Ratio
Body mass index	-0.107	0.045	(-0.182, -0.019)	0.90
Dialyze center <sup>a</sup> (Ref: Center No. 1)				
Dialyze center No. 3	-0.167	0.047	(-2.824, -0.013)	0.85
Level of education (Ref: Illiterate)				
Diploma	-0.673	0.021	(-4.089, -0.563)	0.51
Job (Ref: Employee)				
Unemployed	0.507	0.004	(0.411, 3.250)	1.66
Farmer	0.252	0.005	(0.129, 5.624)	1.29
Number of dialysis sessions per week	-0.421	0.022	(-3.273, -0.202)	0.66
Duration of dialysis sessions	0.802	0.010	(0.341, 3.301)	2.23

Note. CI: Confidence interval.

<sup>a</sup> The name of the centers has been omitted to observe ethics.

respectively. This result may be related to economic issues so that having a job and income plays a more effective role in people feeding, caring, and knowledge. In addition, this issue allows them to have a better quality of life and helps them to adhere to treatment. Given the increased cost of treatment, employees are more successful in meeting their therapeutic needs (22, 24-26).

Additionally, the results showed that with an increase in the frequency of dialysis per week, the risk of death was reduced by 0.34. In other words, increasing the frequency of dialysis leads to increased appetite and food intake and thus improves nutrition. It also has important cardiovascular properties in hemodialysis patients, including lowering blood pressure, reducing environmental resistance, and increasing clearance or reducing endothelial toxins. This justification is also consistent with the findings of several studies (27-29). However, the number of dialysis sessions does not always follow the mentioned justification. In fact, non-adherence to treatment and subsequent electrolyte disturbances in some cases lead to an increase in dialysis sessions. Such a case has been mentioned in some studies (30, 31).

The results further revealed that one unit increase in the duration of hemodialysis was associated with an increased risk of death of 2.23 times. This relationship has been reported in other studies (32, 33). The recent result is probably contrary to our expectations and represents the patient's severe condition; although the duration of hemodialysis is increased, the person is at increased risk of death. It can be explained by the fact that increasing the frequency of hemodialysis increases the risk of hepatitis C virus infection because this infection will be associated with blood transfusions on hemodialysis and can be reduced as much as possible by injecting erythropoietin (34). Even the duration of hemodialysis will be directly related to hepatitis C infection, and the longer the duration of hemodialysis treatment, the greater the need for blood and subsequent hepatitis C infection (35, 36).

### Limitations

This study has some limitations. Compared to some other studies, the sample size is small in this study, and we did not have a sample size. We had a census in all hospitals in Bandar Abbas. Thus, the results of this study cannot be generalized to other cities of Iran. The death rate in our study was low; the medical record departments of the hospitals could not supply all patient information, thus we changed our analysis by the Lasso regression method in the Cox proportional hazard model.

### Conclusion

Overall, education level, job, frequency of hemodialysis per week, and duration of hemodialysis per hour were the most important parameters contributing to the survival of hemodialysis patients. Accordingly, it seems that more education for nurses working in hemodialysis wards and the involved patients can be useful for heeding the recommendations of physicians regarding the required duration for hemodialysis per hour and the number of sessions per week.

### Acknowledgment

The authors would like to thank the department managers of dialysis in the three hospitals of Bandar Abbas and the patients who participated in this study.

### Author Contributions

All authors have contributed to this manuscript and approved this version for submission. LH, SHR, and MT designed the study. LH and AS contributed to the recruitment of patients and the collection of data. In addition, LH and SHR processed and analyzed the results. LH, SHR, MT, and AS wrote the first draft of the manuscript. LH, SHR, and MT helped in revising the manuscript. All authors critically reviewed the manuscript.

### Availability of Data and Materials

The datasets used and/or analyzed during the current study are available from the corresponding author upon reasonable request.

### Competing Interests

The authors declared that they had no competing interests.



**Consent for Publication**

Not applicable.

**Ethical Approval**

The approval was obtained from the Ethics Committee of Hormozgan University of Medical Sciences (Approval number: HUMS.REC.1394.171). Written and informed consent forms were taken from patients to participate in the study. No experimental medical intervention or experimental treatment was used for the particular clinical setting.

**Funding**

None.

**References**

- González R, López-Herce J. Mortality of patients with chronic disease: an increasing problem. *J Pediatr (Rio J)*. 2019;95(6):625-7. doi: [10.1016/j.jpmed.2018.12.002](https://doi.org/10.1016/j.jpmed.2018.12.002).
- Raghupathi W, Raghupathi V. An empirical study of chronic diseases in the United States: a visual analytics approach. *Int J Environ Res Public Health*. 2018;15(3):431. doi: [10.3390/ijerph15030431](https://doi.org/10.3390/ijerph15030431).
- Dehvan F, Monjazebi F, Erfani Khangahi M, Mohammadi H, Ghanei Gheslgh R, Kurdi A. Adequacy of dialysis in Iranian patients undergoing hemodialysis: a systematic review and meta-analysis. *Nephrourol Mon*. 2018;10(5):e82235. doi: [10.5812/numonthly.82235](https://doi.org/10.5812/numonthly.82235).
- Nobakht Haghighi A, Broumand B, D'Amico M, Locatelli F, Ritz E. The epidemiology of end-stage renal disease in Iran in an international perspective. *Nephrol Dial Transplant*. 2002;17(1):28-32. doi: [10.1093/ndt/17.1.28](https://doi.org/10.1093/ndt/17.1.28).
- Ghiasi B, Sarokhani D, Hasanpour Dehkordi A, Sayehmiri K, Heidari MH. Quality of life of patients with chronic kidney disease in Iran: systematic review and meta-analysis. *Indian J Palliat Care*. 2018;24(1):104-11. doi: [10.4103/ijpc.ijpc\\_146\\_17](https://doi.org/10.4103/ijpc.ijpc_146_17).
- Brunner LS. *Brunner & Suddarth's Textbook of Medical-Surgical Nursing*. Vol 1. Lippincott Williams & Wilkins; 2010.
- Raiesifar A, Torabpur M, Mohsenizad P. Causes of chronic renal failure in hemodialysis patients of Abadan. *Iran J Crit Care Nurs*. 2010;2(4):157-60. (Persian).
- Burtis CA, Ashwood ER, Bruns DE. *Tietz Fundamentals of Clinical Chemistry and Molecular Diagnostics-E-Book*. Elsevier Health Sciences; 2012.
- Fanali G, di Masi A, Trezza V, Marino M, Fasano M, Ascenzi P. Human serum albumin: from bench to bedside. *Mol Aspects Med*. 2012;33(3):209-90. doi: [10.1016/j.mam.2011.12.002](https://doi.org/10.1016/j.mam.2011.12.002).
- Gilani N, Kazemnejad A, Zayeri F. Determining the factors affecting albumin serum in hemodialysis patients using random effects model. *J Mazandaran Univ Med Sci*. 2015;25(131):40-7. (Persian).
- Bashardoust B, Gavami M, Maleki N, Doustkami H, Habibzadeh S. Mortality and nutritional status in patients undergoing hemodialysis. *Shiraz E Med J*. 2015;16(2):e20076. doi: [10.17795/semj20076](https://doi.org/10.17795/semj20076).
- Hasan LM, Shaheen DAH, El Kannishy GAH, Sayed-Ahmed NAH, Abd El Wahab AM. Is health-related quality of life associated with adequacy of hemodialysis in chronic kidney disease patients? *BMC Nephrol*. 2021;22(1):334. doi: [10.1186/s12882-021-02539-z](https://doi.org/10.1186/s12882-021-02539-z).
- Nafar M, Samavat S, Khoshdel A, Alipour Abedi B. Dialysis adequacy, dialyzer clearance, and strategies to achieve target: a nationwide multicenter study. *Nephrourol Mon*. 2017;9(1):e42769. doi: [10.5812/numonthly.42769](https://doi.org/10.5812/numonthly.42769).
- Biniaz V, Karimi Moonaghi H, Froutan R, Ebadi A. Kt/V: a magical formula for dialysis adequacy: a critical review. *Jundishapur J Chronic Dis Care*. 2017;6(4):e57280. doi: [10.5812/jjcdc.57280](https://doi.org/10.5812/jjcdc.57280).
- Rezaiee O, Shahgholian N, Shahidi S. Assessment of hemodialysis adequacy and its relationship with individual and personal factors. *Iran J Nurs Midwifery Res*. 2016;21(6):577-82. doi: [10.4103/1735-9066.197673](https://doi.org/10.4103/1735-9066.197673).
- Schober P, Vetter TR. Survival analysis and interpretation of time-to-event data: the tortoise and the hare. *Anesth Analg*. 2018;127(3):792-8. doi: [10.1213/ane.0000000000003653](https://doi.org/10.1213/ane.0000000000003653).
- Badve SV, Paul SK, Klein K, Clayton PA, Hawley CM, Brown FG, et al. The association between body mass index and mortality in incident dialysis patients. *PLoS One*. 2014;9(12):e114897. doi: [10.1371/journal.pone.0114897](https://doi.org/10.1371/journal.pone.0114897).
- Pahlevani V, Fallahzadeh H, Pahlevani N, Nikpour A, Mohammadzadeh M. The effect of time-dependent prognostic factors on survival of non-small cell lung cancer using Bayesian extended cox model. *Iran J Health Sci*. 2020;8(3):29-36. doi: [10.18502/jhs.v8i3.4311](https://doi.org/10.18502/jhs.v8i3.4311).
- Schroijen MA, van de Luijngaarden MW, Noordzij M, Ravani P, Jarraya F, Collart F, et al. Survival in dialysis patients is different between patients with diabetes as primary renal disease and patients with diabetes as a co-morbid condition. *Diabetologia*. 2013;56(9):1949-57. doi: [10.1007/s00125-013-2966-1](https://doi.org/10.1007/s00125-013-2966-1).
- Andrianesis V, Doupis J. The role of kidney in glucose homeostasis--SGLT2 inhibitors, a new approach in diabetes treatment. *Expert Rev Clin Pharmacol*. 2013;6(5):519-39. doi: [10.1586/17512433.2013.827399](https://doi.org/10.1586/17512433.2013.827399).
- Rafiee Vardanjani L, Parvin N, Mahmoodi Shan GR, Molaie E, Shariati A, Hashemina MA. Adherence to hemodialysis treatment and some related factors in hemodialysis patients admitted in Shahrekord Hajar hospital. *J Clin Nurs Midwifery*. 2014;2(4):17-25. (Persian).
- Germin-Petrović D, Mesaros-Devčić I, Lesac A, Mandić M, Soldatić M, Vezmar D, et al. Health-related quality of life in the patients on maintenance hemodialysis: the analysis of demographic and clinical factors. *Coll Antropol*. 2011;35(3):687-93.
- Kalantar-Zadeh K, Kopple JD, Block G, Humphreys MH. A malnutrition-inflammation score is correlated with morbidity and mortality in maintenance hemodialysis patients. *Am J Kidney Dis*. 2001;38(6):1251-63. doi: [10.1053/ajkd.2001.29222](https://doi.org/10.1053/ajkd.2001.29222).
- Borzou SR, Zonoori S, Falahinia GH, Soltanian AR. The effect of education of health-promoting behaviors on lifestyle in hemodialysis patients. *Int J Med Res Health Sci*. 2016;5(6):214-20.
- Masoudi R, Lotfizade M, Gheysarieha MR, Rabiei L. Evaluating the effect of Pender's health promotion model on self-efficacy and treatment adherence behaviors among patients undergoing hemodialysis. *J Educ Health Promot*. 2020;9:197. doi: [10.4103/jehp.jehp\\_747\\_19](https://doi.org/10.4103/jehp.jehp_747_19).
- Hong WP, Lee YJ. The association of dialysis adequacy, body mass index, and mortality among hemodialysis patients. *BMC Nephrol*. 2019;20(1):382. doi: [10.1186/s12882-019-1570-0](https://doi.org/10.1186/s12882-019-1570-0).
- Ahmadmehrabi S, Tang WHW. Hemodialysis-induced cardiovascular disease. *Semin Dial*. 2018;31(3):258-67. doi: [10.1111/sdi.12694](https://doi.org/10.1111/sdi.12694).
- Shafiee MA, Chamanian P, Shaker P, Shahideh Y, Broumand B. The impact of hemodialysis frequency and duration on blood pressure management and quality of life in end-stage renal disease patients. *Healthcare (Basel)*. 2017;5(3):52. doi: [10.3390/healthcare5030052](https://doi.org/10.3390/healthcare5030052).
- Flythe JE, Chang TI, Gallagher MP, Lindley E, Madero M, Sarafidis PA, et al. Blood pressure and volume management in dialysis: conclusions from a Kidney Disease: Improving

- Global Outcomes (KDIGO) Controversies Conference. *Kidney Int.* 2020;97(5):861-76. doi: [10.1016/j.kint.2020.01.046](https://doi.org/10.1016/j.kint.2020.01.046).
30. Ahrari S, Moshki M, Bahrami M. The relationship between social support and adherence of dietary and fluids restrictions among hemodialysis patients in Iran. *J Caring Sci.* 2014;3(1):11-9. doi: [10.5681/jcs.2014.002](https://doi.org/10.5681/jcs.2014.002).
  31. Halle MP, Nelson M, Kaze FF, Jean Pierre NM, Denis T, Fouda H, et al. Non-adherence to hemodialysis regimens among patients on maintenance hemodialysis in sub-Saharan Africa: an example from Cameroon. *Ren Fail.* 2020;42(1):1022-8. doi: [10.1080/0886022x.2020.1826965](https://doi.org/10.1080/0886022x.2020.1826965).
  32. Fotheringham J, Sajjad A, Stel VS, McCullough K, Karaboyas A, Wilkie M, et al. The association between longer haemodialysis treatment times and hospitalization and mortality after the two-day break in individuals receiving three times a week haemodialysis. *Nephrol Dial Transplant.* 2019;34(9):1577-84. doi: [10.1093/ndt/gfz007](https://doi.org/10.1093/ndt/gfz007).
  33. Daugirdas JT. Dialysis time, survival, and dose-targeting bias. *Kidney Int.* 2013;83(1):9-13. doi: [10.1038/ki.2012.365](https://doi.org/10.1038/ki.2012.365).
  34. Beladi Mousavi SS, Shahbazian H, Faramarzi M. Risk factors for developments of hepatitis C and its treatment in patients with end-stage renal disease treated by hemodialysis. *Jundishapur Sci Med J.* 2015;14(1):105-16. (Persian).
  35. Duong CM, Olszyna DP, McLaws ML. Hepatitis B and C virus infections among patients with end stage renal disease in a low-resourced hemodialysis center in Vietnam: a cross-sectional study. *BMC Public Health.* 2015;15:192. doi: [10.1186/s12889-015-1532-9](https://doi.org/10.1186/s12889-015-1532-9).
  36. Fabrizi F, Cerutti R, Messa P. Updated evidence on the epidemiology of hepatitis C virus in hemodialysis. *Pathogens.* 2021;10(9):1149. doi: [10.3390/pathogens10091149](https://doi.org/10.3390/pathogens10091149).